


<p>(51) International Patent Classification ⁶ : C07C 233/49, G01N 33/53, C07C 259/10, C07F 5/02</p>	<p>A1</p>	<p>(11) International Publication Number: WO 98/05627 (43) International Publication Date: 12 February 1998 (12.02.98)</p>
--	-----------	--

Published
With international search report.



Boron compound complexing reagents, intermediate reagents of those reagents and methods of synthesizing these reagents are disclosed. These reagents, including those shown as General Formula (I) and General Formula (II) may be used, after further reactions described herein, to complex with boronic compounds, such as phenylboronic acid or derivatives thereof. Boron compound complexing reagents, boron compound complexes, and methods of synthesizing these reagents and complexes are disclosed. These reagents and complexes include those shown in General Formula (CIII), General formula (CIV), and General Formula (CVI). In one embodiment, the reagents of General formula (CIII) may be used to produce, after condensation with a bioactive species (BAS), the reagent of General Formula (CIV). The reagent of General Formula (CIV) may be used to form a complex with a boron compound, such as a complex shown in General Formula (CVI).

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

-1-

BORONIC COMPOUND COMPLEXING REAGENTS AND HIGHLY STABLE COMPLEXES

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation in part of U.S. application Serial No. 08/188,531 which was filed on January 28, 1994 and is now U.S. Patent No. _____ and which is hereby incorporated herein by reference. This application also follows applications USSNs 08/188,460, 08/188,958 and 08/188, 176, each of which was filed on January 28, 1994 and each of which is incorporated herein by reference.

FIELD OF THE INVENTION

The present invention relates to the field of bioconjugate preparation, and more particularly, to a class of boronic compound complexing reagents useful for the conjugation of biological macromolecules, and the method of making and using such reagents.

BACKGROUND OF THE INVENTION

Bioconjugation is a descriptive term for the joining of two or more different molecular species by chemical or biological means, in which at least one of the molecular species is a biological macromolecule. This includes, but is not limited to, conjugation of proteins, peptides, polysaccharides, hormones, nucleic acids, liposomes and cells, with each other or with any other molecular species that add useful properties, including, but not limited to, drugs, radionuclides, toxins, haptens, inhibitors, chromophores, fluorophores, ligands, etc. Immobilization of biological macromolecules is also considered a special case of bioconjugation in which the macromolecule is conjugated, either reversibly or irreversibly, to an insoluble support. Bioconjugation is utilized extensively in biochemical, immunochemical and molecular biological research. Major applications of bioconjugation include; detection of gene probes, enzyme-linked immuno solid-phase assay, monoclonal antibody drug targeting and medical imaging.

Bioconjugates are generally classified as either direct or indirect conjugates. Direct conjugates encompass those in which two or more components

-2-

are joined by direct covalent chemical linkages. Alternatively, indirect conjugates encompass those in which two or more components are joined via an intermediary complex involving a biological macromolecule. The system described herein is the first to enable the formation of indirect conjugates without dependence upon an intermediary biological macromolecule.

AVIDIN-BIOTIN SYSTEM

Although numerous methods of indirect bioconjugate preparation have been described, a significant number of those reported in the literature have been prepared by exploiting the Avidin-Biotin system, in which, the binding specificity of the protein Avidin (purified from egg white), or Streptavidin (purified from the bacterium *Streptomyces avidinii*), toward the cofactor Biotin (vitamin H) is utilized to bridge an Avidin conjugated macromolecule with a biotinylated macromolecule. Both Avidin and Streptavidin possess four Biotin binding sites of very high affinity ($K = 10^{15} \text{ mol}^{-1}$).

The Avidin-Biotin system has been utilized extensively for enzyme-linked immuno solid-phase assay (ELISA), in which an enzyme-Avidin conjugate (useful for detection by reaction with the enzyme's substrate to afford a colored or chemiluminescent product) is employed to detect the presence of a biotinylated antibody, after first binding the antibody to an immobilized antigen or hapten. Applications of the Avidin-Biotin system number in the hundreds, and have recently been reviewed (Wilchek, M. and Bayer, E. A., (1990) *Methods in Enzymology*, 184).

Although utilized extensively, several limitations are known to be associated with the Avidin-Biotin system, which include nonspecific binding generally attributed to the basicity of the Avidin molecule, nonspecific binding attributed to the presence of carbohydrate residues on the Avidin molecule, and background interference associated with the presence of endogenous Biotin, which is ubiquitous in both eukaryotic and prokaryotic cells.

DIGOXIGENIN- α -DIGOXIGENIN SYSTEM

An alternative indirect bioconjugation system designed to overcome some of the limitations associated with the Avidin-Biotin system has recently been developed for the detection of gene probes by ELISA (Kessler, C., Hölte, H.-J.,

-3-

Seibl, R., Burg, J. and Mühlegger, K., (1990) *Biol. Chem. Hoppe-Seyler*, **371**, 917-965. This system involves the use of the steroid hapten Digoxigenin, an alkaloid occurring exclusively in Digitalis plants, and Fab fragments derived from polyclonal sheep antibodies against Digoxigenin (α -Digoxigenin). The high specificity of the various α -Digoxigenin antibodies affords low backgrounds and eliminates the non-specific binding observed in Avidin-Biotin systems. Digoxigenin-labeled DNA and RNA probes can detect single-copy sequences in human genomic Southern blots. The development of the Digoxigenin α -Digoxigenin system has recently been reviewed (Kessler, C. (1990) in *Advances in Mutagenesis Research* (Obe, G. ed.) pp. 105-152, Springer-Verlag, Berlin/Heidelberg). The Digoxigenin α -Digoxigenin system is the most recent representative of several hapten-antibody systems now utilized for bioconjugation.

IMMOBILIZED PHENYLBORONATES

Phenylboronic acids are known to interact with a wide range of polar molecules having certain requisite functionalities. Complexes of varying stability, involving 1,2-diols, 1,3-diols, 1,2-hydroxy acids, 1,3-hydroxy acids, 1,2-hydroxylamines, 1,3-hydroxylamines, 1,2-diketones and 1,3-diketones, are known to form with either neutral phenylboronic acid or phenylboronate anion. Consequently, immobilized phenylboronic acids have been exploited as chromatographic supports to selectively retain, from diverse biological samples, those molecular species having the requisite functionalities. Many important biological molecules including carbohydrates, catecholamines, prostaglandins, ribonucleosides, and steroids contain the requisite functionalities, and have been either analyzed or purified in this manner. The use of phenylboronic acid chromatographic media for the isolation and separation of biological molecules has been discussed in several reviews (Singhal, R. P. and DeSilva, S. S. M. (1992) *Adv. Chromatog.*, **31**, 293-335; Mazzeo, J. R. and Krull, I. S. (1989) *BioChromatog.*, **4**, 124-130; and Bergold, A. and Scouten, W. H. (1983) in *Solid Phase Biochemistry* (Scouten, W. H. ed.) pp. 149-187, John Wiley & Sons, New York).

Phenylboronic acid, like boric acid, is a Lewis acid, and ionizes not by direct deprotonation, but by hydration to give the tetrahedral phenylboronate anion ($pK_a = 8.86$). Phenylboronic acid is three times as strong an acid as boric acid.

-4-

Ionization of phenylboronic acid is an important factor in complex formation, in that, upon ionization, boron changes from trigonal coordination (having average bond angles of 120° and average bond lengths of 1.37 \AA) to the tetrahedrally coordinated anion (having average bond angles of 109° and average bond lengths of 1.48 \AA).

Molecular species having cis or coaxial 1,2-diol and 1,3-diol functionalities, and particularly carbohydrates, are known to complex with immobilized phenylboronate anion, to form cyclic esters under alkaline aqueous conditions (Lorand, J. P. and Edwards, J. O. (1959) *J. Org. Chem.*, **24**, 769).

Acidification of 1,2-diol and 1,3-diol complexes to neutral pH is known to release the diol containing species, presumably due to hydrolysis of the cyclic ester. Coplanar aromatic 1,3-diols, like 1,8-dihydroxynaphthalene, are known to complex even under acidic conditions due to the hydrolytic stability of six-membered cyclic boronic acid esters (Sienkiewicz, P. A. and Roberts, D. C. (1980) *J. Inorg. Nucl. Chem.*, **42**, 1559-1571). Molecular species having pendant 1,2-hydroxylamine, 1,3-hydroxylamine, 1,2-hydroxyamide, 1,3-hydroxyamide, 1,2-hydroxy-oxime and 1,3-hydroxyoxime functionalities are also known to reversibly complex with phenylboronic acid under alkaline aqueous conditions similar to those associated with the retention of diol containing species (Tanner, D. W. and Bruice, T. C. (1967) *J. Amer. Chem. Soc.*, **89**, 6954).

PHENYLBORONATE BIOCONJUGATES

Ortho-substituted acetamidophenylboronic acids have been proposed as potential linkers for selective bioconjugation via the vicinal diol moieties of the carbohydrate residues associated with glycoproteins (Cai, S. X. and Keana, J. F. W. (1991) *Bioconjugate Chem.*, **2**, 317-322). Phenylboronic acid bioconjugates derived from 3-isothiocyantophenylboronic acid have been successfully utilized for appending radioactive technetium dioxime complexes to monoclonal antibodies for use in medical imaging (Linder, K. E., Wen, M. D., Nowotnik, D. P., Malley, M. F., Gougoutas, J. Z., Nunn, A. D. and Eckelman, W. C. (1991) *Bioconjugate Chem.*, **2**, 160-170; Linder, K. E., Wen, M. D., Nowotnik, D. P., Ramalingam, K., Sharkey, R. M., Yost, F., Narra, R. K. and Eckelman, W. C. (1991) *Bioconjugate Chem.*, **2**, 407-414).

-5-

3-Aminophenylboronic acid has been covalently appended to proteins by a variety of chemical methods and the resulting phenylboronic acid bioconjugates tested for their binding of D-sorbitol, D-mannose and glycated hemoglobin (GHb). The interactions proved to be reversible and of very low affinity rendering the bioconjugates of very limited practical use. Similarly, an alkaline phosphatase phenylboronic acid bioconjugate used in an attempted enzyme-linked assay for the detection of GHb failed to detect the presence of glycated protein (Frantzen, F., Grimsrud, K., Heggli, D. and Sundrehagen, E. (1995) *Journal of Chromatography B*, 670, 37-45).

Although immobilized phenylboronates have been utilized for chromatographic separation of biological molecules having the requisite functionalities, notwithstanding the substantial amount of research into bioconjugation, and the substantial amount of investment in this field, the selectivity of phenylboronic acid has not heretofore been successfully exploited to enable the conjugation of biological macromolecules with one another or with other molecular species that add useful properties.


SUMMARY OF THE INVENTION

The present invention relates to a novel class of boron compound complexing reagents useful for the preparation of bioconjugates, and the method of making and using such reagents. In one embodiment, the boron compound is phenylboronic acid, or derivatives thereof, which complex with the complexing reagents of the present invention. Unless otherwise noted, the phrase phenylboronic acid complexing reagent is used herein to include the broader class of boron compound complexing reagents, and the phrase phenylboronic acid is used herein to include the broader class of boron compounds which complex with the boron compound complexing reagents. In the present invention, in the place of prior art Avidin-Biotin and Digoxigenin α -Digoxigenin systems, boron compound complexing reagents are utilized in conjunction with the boron compound, such as phenylboronic acid reagents (many of which are known in the prior art) to facilitate chemical conjugation without the use of intermediary biological macromolecules. Bioconjugate preparation often involves the conjugation of several components including, but not limited to, proteins, peptides, polysaccharides, hormones, nucleic acids, liposomes and cells, with

-6-

each other or with any other molecular species that add useful properties, including, but not limited to, drugs, radionuclides, toxins, haptens, inhibitors, fluorophores, ligands, solid-phase supports, and boron compound complexing reagents conjugates. These various components utilized in bioconjugate preparation will collectively and individually be termed biologically active species or bioactive species.


Reagents suitable for the modification of a bioactive species for the purpose of incorporating a phenylboronic acid complexing moiety for subsequent conjugation to a different (or the same) bioactive species having pendant phenylboronic acid moieties are of the general formula of General Formula CI:



General Formula CI

Group R_1 is a reactive electrophilic or nucleophilic moiety suitable for reaction of the putative phenylboronic acid complexing reagent with a bioactive species. Group Z is a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide or disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length. Group R_2 is selected from an alkyl (e.g., methyl, ethyl, etc.) and a methylene bearing an electronegative substituent.

Reaction of a reagent of General Formula CI with a bioactive species affords a conjugate having pendant putative phenylboronic acid complexing moieties (one or more) of General Formula CII,




General Formula CII

-7-

The symbol labeled *BAS* represents a biologically active species (or bioactive species) that may or may not contain a portion of a reactive moiety (which may itself have a spacer) used to attach the bioactive species to the reagent. It will be appreciated that, in many embodiments, several identical reagents of the general formula of General Formula CI will react with a single *BAS* molecule. For example, if the *BAS* is a protein, many phenylboronic acid complexing reagents will react with the protein, each reacting at one of the several sites on the protein which are reactive with the R_1 group. Group Z in General Formula CII is a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide or disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length. Group R_2 is selected from an alkyl (e.g., methyl, ethyl, etc.) and a methylene bearing an electronegative substituent.

The reagent of General Formula CII may be further reacted to afford a class of conjugate of boronic compound complexing reagents, e.g., reagents with one or more pendant phenylboronic acid complexing moieties of General Formula CIV:



General Formula CIV


With respect to the conjugate of General Formula CIV, group Z is a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide or ~~disulfide moieties~~, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length. Group R_3 is selected from one of an H, an alkyl, and a methylene or ethylene with an electronegative substituent. Finally, the symbol

-8-

BAS represents the bioactive species that may or may not contain a portion of a reactive moiety used to attach the bioactive species.

The compositions of General Formula CIII and General Formula CIV, and methods for their preparation are described herein and are the subject of my copending application, titled Boronic Compound Complexing Reagents and Highly Stable Complexes, USSN _____, and filed August 5, 1996 by Mark L. Stolowitz, Robert J. Kaiser, and Kevin P. Lund which copending application is hereby incorporated herein by reference.

Phenylboronic acid reagents, many of which are known in the prior art, as well as those described in greater detail in my copending application, titled Phenylboronic Acid Complexes for Bioconjugate Preparation, and filed on January 28, 1994, Serial No. 08/188,958, which is incorporated herein by reference, may be appended to a biologically active species to afford a conjugate having pendant phenylboronic acid moieties (one or more) of General Formula CV:




General Formula CV

wherein the symbol labeled *BAS** represents a second bioactive species, that may include a linker portion and that may differ from the bioactive species labeled *BAS*. The *BAS** may also include a portion of a reactive moiety used to attach the bioactive species to the phenylboronic acid reagent.

A conjugate of General Formula CIV, with at least one biologically active species and having pendent phenylboronic acid complexing moieties (one or more), may be complexed with a conjugate of General Formula CV, prepared from a second bioactive species *BAS** and having pendant phenylboronic acid moieties (one or more), to afford a bioconjugate of General Formula CVI,

-9-



General Formula CVI

wherein the symbols labeled *BAS* and *BAS**, and groups *Z* and *R*₃ are as were previously defined. In this manner, biological macromolecules may be conjugated to one another or with other functionalities which impart useful properties.

Bioconjugates of General Formula CVI may be prepared in buffered aqueous solution or organic solvents. The bioconjugate is formed within a few minutes over a range of temperatures of from about 4° C to 70° C. The stability of the bioconjugate in aqueous solution at a given pH and temperature is significantly influenced by substituent group *R*₃. Bioconjugates of General Formula CVI, are stable in aqueous solutions of approximate pH greater than 3.5 and less than 10.5.


The bioconjugation reaction (phenylboronic acid complexation) is insensitive to significant variations in ionic strength, the presence of organic solvents, the presence of detergents, and the presence of chaotropic agents (protein denaturants), which are incompatible with prior art indirect labeling systems wherein the structure of a biological macromolecule must be maintained to preserve requisite binding properties. In most instances, the constraints governing the formation of bioconjugates, by the system herein described, are limited to those imposed by the conditions required to maintain viability of the bioactive species.

In summary, boron compound complexing reagents, intermediate reagents of those reagents and methods of synthesizing the reagents are described. These reagents, including those shown as General Formulas CI and CII may be used, after further reactions described herein, to complex with boronic compounds, such as phenylboronic acid or derivatives thereof.

Reagents suitable for the modification of a bioactive species for the purpose of incorporating a phenylboronic acid complexing moiety for subsequent

-10-

conjugation to a different (or the same) bioactive species having pendant phenylboronic acid moieties are of the general formula of General Formula CIII:




General Formula CIII

Group R_1 is a reactive electrophilic or nucleophilic moiety suitable for reaction of the phenylboronic acid complexing reagent with a bioactive species. Group Z is a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide or disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length. Group R_3 is selected from one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent.

Group R_3 is preferably selected from one of H, CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 , CH_2CH_2OH and CH_2OCH_3 . When group R_3 is H, group R_1 is preferably selected from one of acrylamide, amino, dithiopyridyl, hydrazide, imidate ester, maleimide, and thiol moieties. Group Z is preferably an unbranched alkyl chain of the general formula $(CH_2)_n$, wherein $n = 1$ to 6.

Reaction of a reagent of General Formula CIII with a bioactive species affords a conjugate having pendant phenylboronic acid complexing moieties (one or more) of General Formula CIV,




General Formula CIV

-11-

The symbol labeled *BAS* represents a biologically active species (or bioactive species) that may or may not contain a portion of a reactive moiety (which may itself have a spacer) used to attach the bioactive species to the reagent. It will be appreciated that, in many embodiments, several identical reagents of the general formula of General Formula CIII will react with a single *BAS* molecule. For example, if the *BAS* is a protein, many phenylboronic acid complexing reagents will react with the protein, each reacting at one of the several sites on the protein which are reactive with the R_1 group. Group *Z* in General Formula CIV is a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide or disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length. Group R_3 is selected from one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent.

The compositions of General Formulas CI and CII, and methods for their preparation are described herein and are the subject of my copending application, titled Boronic Compound Complexing Reagents and Highly Stable Complexes, USSN _____, and filed on August 5, 1996 by Mark L. Stolowitz, Robert J. Kaiser, and Kevin P. Lund, which copending application is hereby incorporated herein by reference.

Phenylboronic acid reagents, many of which are known in the prior art, as well as those described in greater detail in my copending application, titled Phenylboronic Acid Complexes For Bioconjugate Preparation, and filed on January 28, 1994, Serial No. 08/188,958, which is incorporated herein by reference, may be appended to a biologically active species to afford a conjugate having pendant phenylboronic acid moieties (one or more) of General Formula CV:




General Formula CV

wherein the symbol labeled *BAS** represents a second bioactive species, that may include a linker portion and that may differ from the bioactive species labeled

-12-

BAS. The *BAS** may also include a portion of a reactive moiety used to attach the bioactive species to the phenylboronic acid reagent.

A conjugate of General Formula CIV, with at least one biologically active species and having pendent phenylboronic acid complexing moieties (one or more), may be complexed with a conjugate of General Formula CV, prepared from a second bioactive species *BAS** and having pendant phenylboronic acid moieties (one or more), to afford a bioconjugate of General Formula CVI,



General Formula CVI

wherein the symbols labeled *BAS* and *BAS**, and groups *Z* and *R*₃ are as were previously defined. In this manner, as summarized in *Figure 1'*, biological macromolecules may be conjugated to one another or with other functionalities which impart useful properties.

Bioconjugates of General Formula CVI may be prepared in buffered aqueous solution or organic solvents. The bioconjugate is formed within a few minutes over a range of temperatures of from about 4° C to 70° C. The stability of the bioconjugate in aqueous solution at a given pH and temperature is significantly influenced by substituent group *R*₃. Bioconjugates of General Formula CVI, are stable in aqueous solutions of approximate pH greater than 2.5 and less than 12.5. The bioconjugation reaction (phenylboronic acid complexation) is insensitive to significant variations in ionic strength, the presence of organic solvents, the presence of detergents, and the presence of chaotropic agents (protein denaturants), which are incompatible with prior art indirect labeling systems wherein the structure of a biological macromolecule must be maintained to preserve requisite binding properties. In most instances, the constraints governing the formation of bioconjugates, by the system herein described, are limited to those imposed by the conditions required to maintain viability of the bioactive species.

-13-

In summary, boron compound complexing reagents, intermediate reagents of these reagents, boron compound complexes, and methods of synthesizing these reagents and complexes are described. These reagents and complexes include those shown in General Formula CIII, CIV and CVI. In one embodiment, the reagent of General Formula CIII may be used to produce, after condensation with a bioactive species (BAS), the reagent of General Formula CIV. The reagent of General Formula CIV may be used to form a complex with a boron compound, such as a complex shown in General Formula CVI.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 illustrates the utilization of putative phenylboronic acid complexing reagents of General Formula I and phenylboronic acid complexing reagents of General Formula III, to prepare conjugates of General Formula IV which may, in turn, be utilized to prepare bioconjugates of General Formula VI.

Figure 2 summarizes the preparation of alkyl 4- and 5-aminomethylsalicylates, synthetic intermediates leading to reagents of General Formula I, wherein R_2 is an alkyl group, e.g., methyl, ethyl, propyl, etc. Alkyl 4- and 5-aminomethylsalicylates are also useful synthetic intermediates leading to reagents of General Formula III.

Figure 3 summarizes the preparation of alkyl 4- and 5-aminomethylsalicylates, synthetic intermediates leading to reagents of General Formula I, wherein R_2 is a methylene group bearing an electronegative moiety, e.g., carboxymethyl, cyanomethyl, methoxymethyl, etc.

Figure 4 summarizes the preparation of 4- and 5-aminomethylsalicylhydroxamic acids, synthetic intermediates leading to reagents of General Formula III, wherein R_3 is one of either an alkyl group or a methylene or ethylene group bearing an electronegative moiety.

Figure 5 summarizes the synthesis of reagents of General Formulas I and III, wherein R_2 is one of either an alkyl group or a methylene group bearing an

-14-

electronegative moiety, and wherein R₁ is selected from either imidazolidine, hydrazide and *N*-hydroxysuccinimidyl ester moieties.

Figure 6 summarizes the synthesis of reagents of General Formulas I and III, wherein R₂ is one of either an alkyl group or a methylene group bearing an electronegative moiety, and wherein R₁ is selected from either bromo, chloro, iodo, maleimide, dithiopyridyl and imidate ester moieties.

Figure 7 summarizes the synthesis of reagents of General Formulas I and III, wherein R₁ is an *N*-hydroxysuccinimidyl ester, and wherein Z an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of either an intermediate amide or disulfide moiety.

Figure 1' illustrates the utilization of putative phenylboronic acid complexing reagents of General Formula CI and phenylboronic acid complexing reagents of General Formula CIII, to prepare conjugates of General Formula CIV which may, in turn, be utilized to prepare bioconjugates of General Formula CVI.

Figure 2' summarizes the preparation of alkyl 4-aminomethyl-2,6-dihydroxybenzoates, synthetic intermediates leading to reagents of General Formula CI, wherein R₂ is an alkyl group, e.g., methyl, ethyl, propyl, etc. Alkyl 4-aminomethyl-2,6-dihydroxybenzoates are also useful synthetic intermediates leading to reagents of General Formula CIII.

Figure 3' summarizes the preparation of alkyl 4-aminomethyl-2,6-dihydroxybenzoates, synthetic intermediates leading to reagents of General Formula CI, wherein R₂ is a methylene group bearing an electronegative moiety, e.g., carboxymethyl, cyanomethyl, methoxymethyl, etc.

Figure 4' summarizes the preparation of 4-aminomethyl-2,6-dihydroxybenzohydroxamic acids, synthetic intermediates leading to reagents of General Formula CIII, wherein R₃ is one of either an alkyl group or a methylene or ethylene group bearing an electronegative moiety.

-15-

Figure 5' summarizes the synthesis of reagents of General Formulas CI and CIII, wherein R_2 is one of either an alkyl group or a methylene group bearing an electronegative moiety, and wherein R_1 is selected from either imidazolidine, hydrazide and *N*-hydroxysuccinimidyl ester moieties.

Figure 6' summarizes the synthesis of reagents of General Formulas CI and CIII, wherein R_2 is one of either an alkyl group or a methylene group bearing an electronegative moiety, and wherein R_1 is selected from either bromo, chloro, iodo, maleimide, dithiopyridyl and imidate ester moieties.

Figure 7' summarizes the synthesis of reagents of General Formulas CI and CIII, wherein R_1 is an *N*-hydroxysuccinimidyl ester, and wherein Z an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of either an intermediate amide or disulfide moiety.

DETAILED DESCRIPTION OF THE INVENTION


PHENYLBORONIC ACID COMPLEXING REAGENTS AND CONJUGATES DERIVED FROM 4- OR 5-AMINOMETHYLSALICYLIC ACID

This section relates to a class of phenylboronic acid complexing reagents and conjugates derived from 4- or 5-aminomethylsalicylic acid and refers to Figures 1 through 7. Reagents and conjugates derived from 4- or 5-aminomethylsalicylic acid are the subject of my copending applications titled Boronic Compound Complexing Reagents and Complexes, filed August 5, 1996, Serial No. _____ and application titled Boronic Compound Complexing Reagents and Complexes, filed August 5, 1996, Serial No. _____ and Attorney Docket Nos. 81741.P005 and 81741.P006, respectively. The discussion of the preparation of these compounds herein is to facilitate the overall understanding of the distinct and novel invention described herein. By presenting the preparation of various compounds from a similar family of complexing

-16-

reagents and conjugates, it is intended that the full scope of the claimed invention is recognized.

A three-step process which utilizes reagents of General Formula I for the preparation of bioconjugates is summarized in *Figure 1*. Initially, a reagent of General Formula I is selected that is comprised of an appropriate reactive electrophilic or nucleophilic group R_1 suitable for reaction with the desired biologically active species.



General Formula I


Group R_1 is a reactive electrophilic or nucleophilic moiety suitable for reaction of the putative phenylboronic acid complexing reagent with a bioactive species. Group R_1 is preferably selected from, but not limited to, acrylamide, bromo, dithiopyridyl, bromoacetamide, hydrazide, *N*-hydroxysuccinimidyl ester, *N*-hydroxysulfosuccinimidyl ester, imidate ester, imidazolidine, iodo, iodoacetamide, maleimide, amino and thiol moieties.

Group Z is a spacer selected from a saturated or unsaturated, preferably unbranched, chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide or disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length. Group Z is preferably selected from an unbranched alkyl chain of general formula $(CH_2)_n$, wherein $n = 1$ to 6.

Group R_2 is selected from an alkyl (e.g., methyl, ethyl, etc.) and a methylene bearing an electronegative substituent. An electronegative substituent is a substituent with a negative dipole moment, e.g., CN, COOH, etc. Group R_2 is preferably selected from one of CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 and CH_2OCH_3 .

-17-


The next step in a three-step process in the preparation of bioconjugates is to condense the appropriate reagent with the bioactive species to yield a conjugate of General Formula II:



General Formula II

In General Formula II, Z and R₂ are as defined above, and BAS represents a biologically active species which may or may not contain a portion of a reactive moiety (and any spacer) used to attach the biologically active species to the reagent.

Next, the conjugate is reacted with a hydroxylamine derivative of the general formula NH₂OR₃, wherein R₃ is selected from either an H, an alkyl (e.g., methyl, ethyl, etc.), or a methylene or ethylene with an electronegative moiety. Suitable hydroxylamine derivatives include, but are not limited to, NH₂OH, NH₂OCH₃, NH₂OCH₂CN, NH₂OCH₂COOH, NH₂OCH₂CONH₂ and NH₂OCH₂CH₂OH. The resulting conjugate has the general formula of General Formula IV:




General Formula IV

In General Formula IV, Groups Z and BAS are as defined above, and R₃ is selected from one of an H, an alkyl, and a methylene moiety with an electronegative substituent.


The conjugate of General Formula IV is then complexed with a phenylboronic acid having the general formula of General Formula V:

-18-




General Formula V

wherein the symbol labeled *BAS** represents a second biologically active species, that may include a linker portion and differ from the biologically active species labeled *BAS* of the complexing reagent. The *BAS** may also include a portion of a reactive moiety used to attach the bioactive species to the phenylboronic acid reagent. The complexation yields the stereoisomeric complex (about tetrahedral boron) of General Formula VI:



General Formula VI

SYNTHESIS OF PUTATIVE PHENYLBORONIC ACID COMPLEXING REAGENTS OF GENERAL FORMULA I



General Formula I

Reagents of General Formula I are derived from either 4- or 5-methylsalicylic acid. In each instance, the reagent is ultimately prepared from a

-19-

synthetic intermediate which is either an alkyl 4- or 5-aminomethylbenzoate. **Figure 2** summarizes the preparation of alkyl 4- and 5-aminomethylbenzoates, synthetic intermediates leading to reagents of General Formula I, wherein R_2 is an alkyl group, e.g., methyl, ethyl, etc. Figure 2 shows an example where R_2 is a methyl group. Initially, in step 1, either 4- or 5-methyl-salicylic acid is esterified to afford the corresponding alkyl 4- or 5-methylsalicylate. In step 2, the ester is brominated with *N*-bromosuccinimide and benzoyl peroxide catalyst to afford the corresponding benzyl bromide. In step 3, the benzyl bromide is alkylated with sodium azide to afford the corresponding benzyl azide. Finally, in step 4, the benzyl azide is subjected to palladium catalyzed hydrogenation in the presence of HCl to afford the corresponding benzyl amine hydrochloride.

Figure 3 summarizes the preparation of synthetic intermediates leading to reagents of General Formula I, wherein group R_2 is a methylene bearing an electronegative moiety, for example, carboxymethyl, acetamidomethyl and cyanomethyl 4- or 5-aminomethylsalicylate. It is to be appreciated that other contemplated substituents for group R_2 may be prepared by a similar synthesis. Initially, in step 1 of **Figure 3**, the alkyl 4- or 5-aminomethylsalicylate, prepared as summarized in **Figure 2**, is reacted with di-*tert*-butyl dicarbonate and triethylamine in methanol to afford the corresponding protected methyl *N*-(*tert*-butoxycarbonyl)aminomethylbenzoate. In step 2, the methyl ester is cleaved by reaction with potassium trimethylsilanolate and worked up in aqueous acid to afford the corresponding benzoic acid. In step 3, the benzoic acid is alkylated by reaction with either an α -haloacid, α -haloacetamide or α -haloacetonitrile and triethylamine to afford the corresponding carboxymethyl, acetamidomethyl or cyanomethyl ester, respectively. Finally, in step 4, the *N*-(*tert*-butoxycarbonyl) protecting group is removed by reaction with anhydrous HCl in tetrahydrofuran to afford the corresponding benzyl amine hydrochloride.

Another reagent which is the acrylic acid amide of aminomethyl-salicylate, can be prepared in a single step, using the end product of **Figure 3** by condensing acrylic acid anhydride or acryloyl chloride with aminomethyl-salicylate.

Figure 5 summarizes the synthesis of reagents of General Formula I, wherein group R_2 is one of either an alkyl or a methylene bearing an electronegative moiety, and wherein group R_1 is selected from either imidazolidine, hydrazide and *N*-hydroxysuccinimidyl ester moieties. These reagents are each

-20-

prepared by a two-step process in which an aliphatic acid anhydride is utilized in the first step. Initially, an alkyl 4- or 5-aminomethylsalicylate, prepared as summarized in either *Figure 2* or *Figure 3*, is condensed with an aliphatic acid anhydride preferably selected from, but not limited to, either succinic anhydride, glutaric anhydride, maleic anhydride and glycolic acid anhydride, in an aprotic organic solvent, which results in the introduction of a spacer (group Z) having a free terminal carboxylic acid moiety. In the case where the aliphatic acid anhydride is maleic anhydride in this condensation reaction, the resulting Z group is unsaturated as it contains an alkene group. Subsequently, the carboxylic acid moiety is further functionalized by reaction with either *N,N*-carbonyldiimidazole, isobutylchloroformate and *tert*-butyl carbazate, or *N,N*-dicyclohexylcarbodiimide and *N*-hydroxysuccinimide to afford the corresponding imidazolidine, protected hydrazide and *N*-hydroxysuccinimidyl ester, respectively. In the instance of the protected hydrazide, the *N*-(*tert*-butoxycarbonyl) protecting group is removed by contacting the reagent with anhydrous hydrochloric acid.

Figure 6 summarizes the synthesis of reagents of General Formula I, wherein group R₂ is one of either an alkyl or a methylene bearing an electronegative moiety, and wherein group R₁ is selected from either acrylamide, bromoacetamide, chloroacetamide, maleimide, dithiopyridyl and imidate ester moieties. Reagents of General Formula I, wherein group R₁ is selected from either bromo and chloro moieties, are prepared by condensing an alkyl 4- or 5-aminomethylsalicylate, prepared as summarized in either *Figure 2* or *Figure 3*, with either bromoacetic acid anhydride or chloroacetic acid, respectively. The homologous iodo reagent is prepared by halogen exchange of the chloro reagent with sodium iodide. Reagents of General Formula I, wherein R₁ is selected from either bromo, chloro, iodo, bromoacetamide, chloroacetamide and iodoacetamide moieties, may not be conveniently prepared when R₃ is H, due to the potential for intermolecular alkylation of the unprotected hydroxamate. Reagents of General Formula I, wherein group R₁ is selected from either maleimide or dithiopyridyl moieties, are prepared by condensing an alkyl 4- or 5-aminomethylsalicylate, prepared as summarized in either *Figure 2* or *Figure 3*, with an *N*-hydroxysuccinimidyl ester of an aliphatic carboxylic ester which bears either a terminal maleimide to dithiopyridyl moiety. A reagent of General Formula I, wherein R₁ is an imidate ester moiety, is prepared by a two-step process in which

-21-

an alkyl 4- or 5-aminomethyl-salicylate, prepared as summarized in either *Figure 2* or *Figure 3*, is first condensed with an *N*-hydroxysuccinimidyl ester of an aliphatic carboxylic ester which bears a terminal nitrile moiety. Subsequently, the nitrile moiety is converted to the methyl imidate ester by reaction with anhydrous hydrogen chloride in methanol at 0° C.

Reagents of General Formula I, wherein group R₁ is selected from either *N*-hydroxy-succinimidyl ester and dithiopyridyl moieties, may be utilized as synthetic intermediates to prepare reagents of General Formula I, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide or disulfide moieties.

Reagents of General Formula I, wherein group R₁ is an *N*-hydroxysuccinimidyl ester moiety, may be condensed with compounds having primary aliphatic amine moieties of the general formula R₁-Z₂-NH₂, wherein Z₂ is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford reagents of General Formula I, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide moieties.

Alternatively, *N*-hydroxysuccinimidyl ester reagents of General Formula I, preferably derived from a dicarboxylic acid selected from either succinic acid, maleic acid, fumaric acid, acetylenedicarboxylic acid and glutaric acid, may be condensed with compounds having primary aliphatic amine moieties of the general formula HO₂C-Z₂-NH₂, wherein Z₂ is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, preferably selected from, but not limited to, either glycine, β-alanine, aminopropionic acid, 4-aminobutyric acid and 6-aminocaproic acid, to afford compounds having free terminal carboxylic acid moieties which may be further functionalized in accordance with *Figure 5* to afford reagents of General Formula I, wherein Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide moieties. This process is summarized in *Figure 7* for the synthesis of a reagent of General Formula I, wherein R₁ is an *N*-hydroxysuccinimidyl ester, and wherein Z is an unbranched saturated or unsaturated chain with at least one of an intermediate amide moiety.

Reagents of General Formula I, wherein group R₁ is a dithiopyridyl moiety, may be condensed with compounds having terminal thiol moieties of the


general formula R_1-Z_2-SH , wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford reagents of General Formula I, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate disulfide moieties.

Alternatively, dithiopyrinyll reagents of General Formula I, preferably derived from a mercaptocarboxylic acid selected from either mercaptoacetic acid, β -mercaptopropionic acid, mercaptopropionic acid, 4-mercaptobutyric acid and 6-mercaptocaproic acid, may be condensed with compounds having a thiol moiety of the general formula HO_2C-Z_2-SH , wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, preferably selected from, either mercaptoacetic acid, β -mercaptopropionic acid, mercaptopropionic acid, 4-mercaptobutyric acid and 6-mercaptocaproic acid, to afford compounds having free terminal carboxylic acid moieties which may be further functionalized in accordance with *Figure 5* to afford reagents of General Formula I, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate disulfide moieties. This process is summarized in *Figure 7* for the synthesis of a reagent of General Formula I, wherein R_1 is an *N*-hydroxysuccinimidyl ester, and wherein Z is an unbranched saturated or unsaturated chain with at least one of an intermediate disulfide moiety.

Reagents of General Formula I, wherein group Z is a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length, are prepared by condensing an alkyl 4- or 5-aminomethylsalicylate, prepared as summarized in Figures 2 or 3, with a polyethylene glycol reagent having both an *N*-hydroxysuccinimidyl ester moiety and either a reactive electrophilic or nucleophilic moiety (or a protected precursor thereof), many of which are commercially available, to afford reagents of General Formula I, wherein group Z is a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length.

-23-

SYNTHESIS OF
PHENYLBORONIC ACID COMPLEXING REAGENTS
OF GENERAL FORMULA III




General Formula III

Figure 4 summarizes the preparation of 4- and 5-aminomethylsalicylhydroxamic acids, synthetic intermediates leading to reagents of General Formula III, wherein group R_3 is one of an alkyl or methylene bearing an electronegative moiety. Initially, in step 1, an alkyl 4- or 5-aminomethylsalicylate, prepared as summarized in **Figure 2**, is condensed with *N*-(benzyloxy-carbonyl)oxy succinimide to afford the alkyl *N*-benzyloxycarbonyl protected 4- or 5-amino-methylsalicylate. In step 2, the phenolic hydroxyl moiety was condensed with benzyl bromide to afford the further protected benzyl ether intermediate. In step 3, the alkyl ester is selectively cleaved by reaction with LiOH to afford the corresponding benzoic acid. In step 4, the benzoic acid is activated by reaction with isobutylchloroformate to form a mixed anhydride which is subsequently reacted with a hydroxylamine derivative preferably selected from, but not limited to, either NH_2OH , NH_2OCH_3 , $\text{NH}_2\text{OCH}_2\text{CN}$, $\text{NH}_2\text{OCH}_2\text{COOH}$, $\text{NH}_2\text{OCH}_2\text{CONH}_2$, $\text{NH}_2\text{OCH}_2\text{CH}_2\text{OH}$ and $\text{NH}_2\text{OCH}_2\text{OCH}_3$ to afford the corresponding protected hydroxamic acid. Finally, in step 5, both the amine and phenolic hydroxyl moieties are simultaneously deprotected by palladium catalyzed hydrogenation in the presence of HCl to afford the corresponding 4- or 5-aminomethylsalicylhydroxamic acid hydrochloride. These synthetic intermediates, which are structurally related to the corresponding alkyl 4- or 5-aminomethylsalicylates, may be further functionalized in accordance with either **Figure 5** or **Figure 6** to afford reagents of General Formula III.

-24-


PREPARATION OF
PHENYLBORONIC ACID COMPLEXING CONJUGATES
OF GENERAL FORMULA IV

At this point, the putative phenylboronic acid complexing reagents of General Formula I may be reacted with a suitable biologically active species to yield the conjugate of the general formula of General Formula II:



General Formula II

The conjugate of General Formula II is next condensed with a hydroxylamine derivative to yield the corresponding phenylboronic acid complexing conjugate of the general formula of General Formula IV:




General Formula IV

Suitable hydroxylamine derivatives are preferably selected from, but not limited to, NH_2OH , NH_2OCH_3 , $\text{NH}_2\text{OCH}_2\text{CN}$, $\text{NH}_2\text{OCH}_2\text{COOH}$, $\text{NH}_2\text{OCH}_2\text{CONH}_2$, $\text{NH}_2\text{OCH}_2\text{CH}_2\text{OH}$, $\text{NH}_2\text{OCH}_2\text{OCH}_3$. When group R_2 in General Formula II is an alkyl group, NH_2OH is preferably utilized to effect the interconversion of General Formula II to General Formula IV.


Alternatively, conjugates of General Formula IV may also be prepared by the route which is described in *Figure 1* which utilizes a reagent of General Formula III:

-25-




General Formula III

Phenylboronic acid complexing reagents of General Formula III may be reacted with a suitable biologically active species to yield the conjugate of the general formula of General Formula IV:



General Formula IV

A two-step process which utilizes reagents of General Formula III for the preparation of bioconjugates is summarized in *Figure 1*. Initially, a reagent of General Formula III is selected that is comprised of an appropriate reactive electrophilic or nucleophilic group R_1 suitable for reaction with the desired biologically active species.



General Formula III

Group R_1 is a reactive electrophilic or nucleophilic moiety suitable for reaction of the phenylboronic acid complexing reagent with a bioactive species. Group R_1 is preferably selected from, but not limited to, acrylamide, bromo, dithiopyridyl, bromoacetamide, hydrazide, *N*-hydroxysuccinimidyl ester, *N*-hydroxysulfosuccinimidyl ester, imidate ester, imidazolidine, iodo, iodoacetamide, maleimide, amino and thiol moieties.


-26-

Group Z is a spacer selected from a saturated or unsaturated, preferably unbranched, chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide or disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length. Group Z is preferably selected from an unbranched alkyl chain of general formula $(CH_2)_n$, wherein $n = 1$ to 6.

Group R_3 is selected from one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent.

Group R_3 is preferably selected from one of H, CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 , CH_2CH_2OH and CH_2OCH_3 . When group R_3 is H, group R_1 is preferably selected from one of acrylamide, amino, dithiopyridyl, hydrazide, imidate ester, maleimide, and thiol moieties.


The reagent of General Formula III is condensed with the bioactive species to yield a conjugate of General Formula IV:



General Formula IV

wherein groups Z and BAS are as defined above, and R_3 is selected from one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent.

The conjugate of General Formula IV is then complexed with a phenylboronic acid conjugate having the general formula of General Formula V:




General Formula V

wherein the symbol labeled BAS^* represents a second biologically active species, that may include a linker portion and differ from the biologically active species


-27-

labeled *BAS* of the complexing reagent. The *BAS** may also include a portion of a reactive moiety used to attach the bioactive species to the phenylboronic acid reagent. The complexation yields the stereoisomeric complex (tetrahedral boron) of General Formula VI:



General Formula VI

An alternative three-step process which utilizes reagents of General Formula I for the preparation of bioconjugates is also summarized in *Figure 1*. Initially, a reagent of General Formula I is selected that is comprised of an appropriate reactive electrophilic or nucleophilic group R_1 suitable for reaction with the desired biologically active species.



General Formula I

Group R_1 is a reactive electrophilic or nucleophilic moiety suitable for reaction of the putative phenylboronic acid complexing reagent with a bioactive species. Group R_1 is preferably selected from, but not limited to, acrylamide, bromo, dithiopyridyl, bromoacetamide, hydrazide, *N*-hydroxysuccinimidyl ester, *N*-hydroxysulfosuccinimidyl ester, imidate ester, imidazolidine, iodo, iodoacetamide, maleimide, amino and thiol moieties.


Group Z is a spacer selected from a saturated or unsaturated, preferably unbranched, chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide or disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length.

-28-

Group Z is preferably selected from an unbranched alkyl chain of general formula $(CH_2)_n$, wherein $n = 1$ to 6.

Group R_2 is selected from an alkyl (e.g., methyl, ethyl, etc.) and a methylene or ethylene bearing an electronegative substituent. An electronegative substituent is a substituent with a negative dipole moment, e.g., CN, COOH, etc. Group R_2 is preferably selected from one of CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 and CH_2OCH_3 .


The next step in a three-step process in the preparation of bioconjugates is to condense the appropriate reagent with the bioactive species to yield a conjugate of General Formula II:



General Formula II

In General Formula II, Z, BAS and R_2 are as defined above.

Next, the conjugate is reacted with a hydroxylamine derivative of the general formula NH_2OR_3 , wherein R_3 is selected from either an H, an alkyl (e.g., methyl, ethyl, etc.), or a methylene or ethylene with an electronegative substituent. Suitable hydroxylamine derivatives include, but not limited to, NH_2OH , NH_2OCH_3 , NH_2OCH_2CN , NH_2OCH_2COOH , $NH_2OCH_2CONH_2$ and $NH_2OCH_2CH_2OH$. The resulting conjugate has the general formula of General Formula IV:




General Formula IV

wherein groups Z and BAS are as defined above, and R_3 is selected from one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent.


-29-

The conjugate of General Formula IV is then complexed with a phenylboronic acid having the general formula of General Formula V:




General Formula V

wherein the symbol labeled *BAS** represents a second biologically active species, that may include a linker portion and differ from the biologically active species labeled *BAS* of the complexing reagent. The *BAS** may also include a portion of a reactive moiety used to attach the bioactive species to the phenylboronic acid reagent. The complexation yields the stereoisomeric complex (tetrahedral boron) of General Formula VI:



General Formula VI

SYNTHESIS OF PHENYLBORONIC ACID COMPLEXING REAGENTS OF GENERAL FORMULA III



General Formula III

-30-

Reagents of General Formula III are derived from either 4- or 5-methylsalicylic acid. In each instance, the reagent is ultimately prepared from a synthetic intermediate which is either an alkyl 4- or 5-aminomethylbenzoate.

Figure 2 summarizes the preparation of alkyl 4- and 5-aminomethylsalicylates, synthetic intermediates leading to reagents of General Formula III, wherein R_2 is an alkyl group, e.g., methyl, ethyl, etc. Figure 2 shows an example where R_2 is a methyl group. Initially, in step 1, either 4- or 5-methylsalicylic acid is esterified to afford the corresponding alkyl 4- or 5-methylsalicylate. In step 2, the ester is brominated with *N*-bromo-succinimide and benzoyl peroxide catalyst to afford the corresponding benzyl bromide. In step 3, the benzyl bromide is alkylated with sodium azide to afford the corresponding benzyl azide. Finally, in step 4, the benzyl azide is subjected to palladium catalyzed hydrogenation in the presence of HCl to afford the corresponding benzyl amine hydrochloride.

Figure 4 summarizes the preparation of 4- and 5-aminomethylsalicylhydroxamic acids, synthetic intermediates leading to reagents of General Formula III, wherein group R_3 is one of an alkyl or methylene or ethylene bearing an electronegative substituent. Initially, in step 1, an alkyl 4- or 5-aminomethylsalicylate, prepared as summarized in **Figure 2**, is condensed with *N*-(benzyloxycarbonyl)oxy succinimide to afford the alkyl *N*-benzyloxycarbonyl protected 4- or 5-aminomethylsalicylate. In step 2, the phenolic hydroxyl moiety was condensed with benzyl bromide to afford the further protected benzyl ether intermediate. In step 3, the alkyl ester is selectively cleaved by reaction with LiOH to afford the corresponding benzoic acid. In step 4, the benzoic acid is activated by reaction with isobutylchloroformate to form a mixed anhydride which is subsequently reacted with a hydroxylamine derivative preferably selected from, but not limited to, either NH_2OH , NH_2OCH_3 , NH_2OCH_2CN , NH_2OCH_2COOH , $NH_2OCH_2CONH_2$, $NH_2OCH_2CH_2OH$ and $NH_2OCH_2OCH_3$ to afford the corresponding protected hydroxamic acid. Finally, in step 5, both the amine and phenolic hydroxyl moieties are simultaneously deprotected by palladium catalyzed hydrogenation in the presence of HCl to afford the corresponding 4- or 5-aminomethylsalicylhydroxamic acid hydrochloride.

Another reagent of the present invention, which is the acrylic acid amide of aminomethyl-salicylhydroxamic acid, can be prepared in a single step, using the

end product of Figure 4 by condensing acrylic acid anhydride or acryloyl chloride with aminomethyl-salicylhydroxamic acid, provided that R_3 is not H.

Figure 5 summarizes the synthesis of reagents of General Formula III, wherein group R_3 is preferably selected from one of H, CH_3 , and a methylene or ethylene moiety with an electronegative substituent, and wherein group R_1 is selected from either imidazolidine, hydrazide and *N*-hydroxysuccinimidyl ester moieties. These reagents are each prepared by a two-step process in which an aliphatic acid anhydride is utilized in the first step. Initially, a 4- or 5-aminomethylsalicylhydroxamic acid, prepared as summarized in **Figure 4**, is condensed with an aliphatic acid anhydride preferably selected from, but not limited to, either succinic anhydride, glutaric anhydride, maleic anhydride and glycolic acid anhydride, in an aprotic organic solvent, which results in the introduction of a spacer (group Z) having a free terminal carboxylic acid moiety. In the case where the aliphatic acid anhydride is maleic anhydride in this condensation reaction, the resulting Z group is unsaturated as it contains an alkene group. Subsequently, the carboxylic acid moiety is further functionalized by reaction with either *N,N*-carbonyl-diimidazole, isobutylchloroformate and *tert*-butyl carbazate, or *N,N*-dicyclohexylcarbodiimide and *N*-hydroxysuccinimide to afford the corresponding imidazolidine, protected hydrazide and *N*-hydroxysuccinimidyl ester, respectively. In the instance of the protected hydrazide, the *N*-(*tert*-butoxycarbonyl) protecting group is removed by contacting the reagent with anhydrous hydrochloric acid.

Figure 6 summarizes the synthesis of reagents of General Formula III, wherein group R_3 is preferably selected from one of H, CH_3 , and a methylene or ethylene moiety with an electronegative substituent, and wherein group R_1 is selected from either bromo, chloro, maleimide, dithiopyridyl and imidate ester moieties. Reagents of General Formula III, wherein group R_1 is selected from either bromo and chloro moieties, are prepared by condensing a 4- or 5-aminomethylsalicylhydroxamic acid, prepared as summarized in **Figure 4**, with either bromoacetic acid anhydride or chloroacetic acid anhydride, respectively. The homologous iodo reagent is prepared by halogen exchange of the chloro reagent with sodium iodide. Reagents of General Formula III, wherein R_1 is selected from either bromo, chloro, iodo, bromoacetamide, chloroacetamide and iodoacetamide moieties, may not be conveniently prepared when R_3 is H, due to the potential for intermolecular alkylation of the unprotected hydroxamate.

-32-

Reagents of General Formula III, wherein group R_1 is selected from either maleimide and dithiopyridyl moieties, are prepared by condensing a 4- or 5-aminomethylsalicylhydroxamic acid, prepared as summarized in *Figure 4*, with an *N*-hydroxysuccinimidyl ester of an aliphatic carboxylic ester which bears either a terminal maleimide or dithiopyridyl moiety. Reagents of General Formula III, wherein R_1 is an imidate ester moiety, are prepared by a two-step process in which a 4- or 5-aminomethylsalicylhydroxamic acid, prepared as summarized in *Figure 4*, is first condensed with an *N*-hydroxysuccinimidyl ester of an aliphatic carboxylic ester which bears a terminal nitrile moiety. Subsequently, the nitrile moiety is converted to the methyl imidate ester by reaction with anhydrous hydrochloric acid in methanol at 0° C.

Reagents of General Formula III, wherein group R_1 is selected from either *N*-hydroxy-succinimidyl ester and dithiopyridyl moieties may be utilized as synthetic intermediates to prepare reagents of General Formula III, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide or disulfide moieties.

Reagents of General Formula III, wherein group R_1 is an *N*-hydroxysuccinimidyl ester moiety, may be condensed with compounds having primary aliphatic amine moieties of the general formula $R_1-Z_2-NH_2$, wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford reagents of General Formula III, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide moieties.

Alternatively, *N*-hydroxysuccinimidyl ester reagents of General Formula III, preferably derived from a dicarboxylic acid selected from either succinic acid, maleic acid, fumaric acid, acetylenedicarboxylic acid and glutaric acid, may be condensed with compounds having primary aliphatic amine moieties of the general formula $HO_2C-Z_2-NH_2$, wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, preferably selected from, either glycine, β -alanine, α -aminopropionic acid, 4-amino-butyric acid and 6-aminocaproic acid, to afford compounds having free terminal carboxylic acid moieties which may be further functionalized in accordance with *Figure 5* to afford reagents of General Formula III, wherein Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide moieties. This process is

summarized in Figure 7 for the synthesis of a reagent of General Formula III, wherein R_1 is an *N*-hydroxysuccinimidyl ester, and wherein Z is an unbranched saturated or unsaturated chain with at least one of an intermediate amide moiety.


Reagents of General Formula III, wherein group R_1 is a dithiopyridyl moiety, may be condensed with compounds having terminal thiol moieties of the general formula R_1-Z_2-SH , wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford reagents of General Formula III, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate disulfide moieties.

Alternatively, dithiopyridyl reagents of General Formula III, preferably derived from a mercaptocarboxylic acid selected from either mercaptoacetic acid, β -mercaptopropionic acid, mercaptopropiolic acid, 4-mercaptoputyric acid and 6-mercaptopcaproic acid, may be condensed with compounds having a terminal thiol moiety of the general formula HO_2C-Z_2-SH , wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, preferably selected from either mercaptoacetic acid, β -mercaptopropionic acid, mercaptopropiolic acid, 4-mercaptoputyric acid and 6-mercaptopcaproic acid, to afford compounds having free terminal carboxylic acid moieties which may be further functionalized in accordance with Figure 5 to afford reagents of General Formula III, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate disulfide moieties. This process is summarized in Figure 7 for the synthesis of a reagent of General Formula III, wherein R_1 is an *N*-hydroxysuccinimidyl ester, and wherein Z is an unbranched saturated or unsaturated chain with at least one of a intermediate disulfide moiety.

Reagents of General Formula III, wherein group Z is a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length, are prepared by condensing a 4-aminomethyl-salicylhydroxamic acid, prepared as summarized in Figure 4, with a polyethylene glycol reagent having both an *N*-hydroxysuccinimidyl ester moiety and either a reactive electrophilic or nucleophilic moiety (or a precursor thereof), many of which are commercially available, to afford reagents of General Formula III, wherein group Z is a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length.

-34-

SYNTHESIS OF
PUTATIVE PHENYLBORONIC ACID COMPLEXING REAGENTS
OF GENERAL FORMULA I



General Formula I

Reagents of General Formula I are derived from either 4- or 5-methylsalicylic acid. In each instance, the reagent is ultimately prepared from a synthetic intermediate which is either an alkyl 4- or 5-aminomethylsalicylate. **Figure 2** summarizes the preparation of alkyl 4- and 5-aminomethylsalicylates, synthetic intermediates leading to reagents of General Formula I, wherein R_2 is an alkyl group, e.g., methyl, ethyl, etc. Figure 2 shows an example where R_2 is a methyl group. Initially, in step 1, either 4- or 5-methylsalicylic acid is esterified to afford the corresponding alkyl 4- or 5-methylsalicylate. In step 2, the ester is brominated with *N*-bromosuccinimide and benzoyl peroxide catalyst to afford the corresponding benzyl bromide. In step 3, the benzyl bromide is alkylated with sodium azide to afford the corresponding benzyl azide. Finally, in step 4, the benzyl azide is subjected to palladium catalyzed hydrogenation in the presence of HCl to afford the corresponding benzyl amine hydrochloride.

Figure 3 summarizes the preparation of synthetic intermediates leading to reagents of General Formula I, wherein group R_2 is a methylene or ethylene bearing an electronegative moiety, for example, carboxymethyl, acetamidomethyl and cyanomethyl 4- or 5-aminomethylsalicylate. It is to be appreciated that other contemplated substituents for group R_2 may be prepared by a similar synthesis. Initially, in step 1 of **Figure 3**, the alkyl 4- or 5-amino-methylsalicylate, prepared as summarized in **Figure 2**, is reacted with *di-tert*-butyl dicarbonate and triethylamine in methanol to afford the corresponding protected methyl *N*-(*tert*-butoxy-carbonyl)aminomethylbenzoate. In step 2, the methyl ester is cleaved by reaction with potassium trimethylsilanolate and worked up in aqueous acid to

-35-

afford the corresponding benzoic acid. In step 3, the benzoic acid is alkylated by reaction with either an α -haloacid, α -haloacetamide or α -haloacetonitrile and triethylamine to afford the corresponding carboxymethyl, acetamidomethyl or cyanomethyl ester, respectively. Finally, in step 4, the *N*-(*tert*-butoxy-carbonyl) protecting group is removed by reaction with anhydrous HCl in tetrahydrofuran to afford the corresponding benzyl amine hydrochloride. In the instance of the protected hydrazide, the *N*-(*tert*-butoxycarbonyl) protecting group is removed by contacting the reagent with anhydrous hydrochloric acid.

Figure 5 summarizes the synthesis of reagents of General Formula I, wherein group R_2 is one of either an alkyl or a methylene or ethylene bearing an electronegative substituent, and wherein group R_1 is selected from either imidazolidine, hydrazide and *N*-hydroxysuccinimidyl ester moieties. These reagents are each prepared by a two-step process in which an aliphatic acid anhydride is utilized in the first step. Initially, an alkyl 4- or 5-aminomethylsalicylate, prepared as summarized in either **Figure 2** or **Figure 3**, is condensed with an aliphatic acid anhydride preferably selected from, but not limited to, either succinic anhydride, glutaric anhydride, Maleic anhydride, and glycolic acid anhydride, in an aprotic organic solvent, which results in the introduction of a spacer (group Z) having a free terminal carboxylic acid moiety. Subsequently, the carboxylic acid moiety is further functionalized by reaction with either *N,N*-carbonyldiimidazole, isobutylchloroformate and *tert*-butyl carbazate, or *N,N*-dicyclohexylcarbodiimide and *N*-hydroxy-succinimide to afford the corresponding imidazolidine, protected hydrazide and *N*-hydroxy-succinimidyl ester, respectively.

Figure 6 summarizes the synthesis of reagents of General Formula I, wherein group R_2 is one of either an alkyl or a methylene or ethylene bearing an electronegative substituent, and wherein group R_1 is selected from either bromo, chloro, maleimide, dithiopyridyl and imidate ester moieties. Reagents of General Formula I, wherein group R_1 is selected from either bromo and chloro moieties, are prepared by condensing an alkyl 4- or 5-aminomethylsalicylate, prepared as summarized in either **Figure 2** or **Figure 3**, with either bromoacetic acid anhydride or chloroacetic acid anhydride, respectively. The homologous iodo reagent is prepared by halogen exchange of the chloro reagent with sodium iodide. Reagents of General Formula III, wherein R_1 is selected from either bromo, chloro, iodo, bromoacetamide, chloroacetamide and iodoacetamide moieties, may

-36-

not be conveniently prepared when R_3 is H, due to the potential for intermolecular alkylation of the unprotected hydroxamate. Reagents of General Formula I, wherein group R_1 is selected from either maleimide or dithiopyridyl moieties, are prepared by condensing an alkyl 4- or 5-aminomethylsalicylate, prepared as summarized in either *Figure 2* or *Figure 3*, with an *N*-hydroxysuccinimidyl ester of an aliphatic carboxylic ester which bears either a terminal maleimide or dithiopyridyl moiety. A reagent of General Formula I, wherein R_1 is an imidate ester moiety, is prepared by a two-step process in which an alkyl 4- or 5-aminomethylsalicylate, prepared as summarized in either *Figure 2* or *Figure 3*, is first condensed with an *N*-hydroxysuccinimidyl ester of an aliphatic carboxylic ester which bears a terminal nitrile moiety. Subsequently, the nitrile moiety is converted to the methyl imidate ester by reaction with anhydrous hydrochloric acid in methanol at 0° C.

Reagents of General Formula I, wherein group R_1 is selected from either dithiopyridyl, imidazolidine and *N*-hydroxysuccinimidyl ester moieties may be utilized as synthetic intermediates to prepare reagents of General Formula I having extended spacers, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide or disulfide moieties.

Reagents of General Formula I, wherein group R_1 is selected from either imidazolidine and *N*-hydroxysuccinimidyl ester moieties, may be condensed with compounds having primary aliphatic amine moieties of the general formula $R_1-Z'-NH_2$, wherein Z' is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford reagents of General Formula I, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide moieties. Note that Z' is shown as Z_2 in *Figure 7*.

Alternatively, reagents of General Formula I, wherein group R_1 is selected from either imidazolidine and *N*-hydroxysuccinimidyl ester moieties, may be condensed with compounds having primary aliphatic amine moieties of the general formula $H_2N-Z'-COOH$, wherein Z' is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, preferably selected from, but not limited to, either glycine, β -alanine, 4-aminobutyric acid and 6-aminocaproic acid, to afford compounds having a free terminal carboxylic acid moiety which may be further functionalized in accordance with *Figure 5* to

-37-

afford reagents of General Formula I, wherein Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide moieties.

Reagents of General Formula I, wherein group R_1 is a dithiopyridyl moiety, may be condensed with compounds having terminal thiol moieties of the general formula $R_1-Z'-SH$, wherein Z' is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford reagents of General Formula I, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate disulfide moieties.


Alternatively, reagents of General Formula I, wherein group R_1 is a dithiopyridyl moiety, may be condensed with compounds having terminal thiol moieties of the general formula $HS-Z'-COOH$, wherein Z' is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford compounds having a free terminal carboxylic acid moiety which may be further functionalized in accordance with *Figure 5* to afford reagents of General Formula I, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate disulfide moieties.

Reagents of General Formula I, wherein group Z is a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length, are prepared by condensing an alkyl 4- or 5-amino-methylsalicylate, prepared as summarized in either *Figure 2* or *Figure 3*, with a polyethylene glycol reagent having both an *N*-hydroxysuccinimidyl ester moiety and either a reactive electrophilic or nucleophilic moiety (or a precursor thereof), many of which are commercially available, to afford reagents of General Formula I, wherein group Z is a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length.

PREPARATION OF PHENYLBORONIC ACID COMPLEXING CONJUGATES OF GENERAL FORMULA IV


At this point, phenylboronic acid complexing reagents of General Formula III may be reacted with a suitable biologically active species to yield the conjugate of the general formula of General Formula IV:

-38-




General Formula IV

Alternatively, the putative phenylboronic acid complexing reagents of General Formula I may be reacted with a suitable biologically active species to yield the conjugate of the general formula of General Formula II:



General Formula II

The conjugate of General Formula II is next condensed with a hydroxylamine derivative to yield the corresponding phenylboronic acid complexing conjugate of the general formula of General Formula IV:



General Formula VI

Suitable hydroxylamine derivatives are preferably selected from, but not limited to, NH_2OH , NH_2OCH_3 , $\text{NH}_2\text{OCH}_2\text{CN}$, $\text{NH}_2\text{OCH}_2\text{COOH}$, $\text{NH}_2\text{OCH}_2\text{CONH}_2$, $\text{NH}_2\text{OCH}_2\text{CH}_2\text{OH}$, $\text{NH}_2\text{OCH}_2\text{OCH}_3$. When group R_2 in General Formula II is an alkyl group, NH_2OH is preferably utilized to effect the interconversion of General Formula II to General Formula IV.

-39-

PREPARATION OF BIOCONJUGATES OF GENERAL FORMULA VI

Bioconjugates of General Formula VI may be prepared in buffered aqueous solutions or organic solvents. Preferred buffers include acetate, citrate, phosphate, carbonate and diglycine. Borate buffers should be avoided due to their ability to complex with the phenylboronic acid complexing moiety. Tris, β -hydroxyamine and β -hydroxyacid buffers should be avoided due to their ability to complex with the phenylboronic acid. The bioconjugate is formed within a few minutes over a range of temperatures of from about 4°C to 70°C. The stability of the bioconjugate in aqueous solution at a given pH and temperature is influenced, to some extent, by substituent group R_3 . Bioconjugates of General Formula VI are stable in aqueous solutions of approximate pH greater than 3.5 and less than 10.5. The bioconjugation reaction (phenylboronic acid complexation) is insensitive to significant variations in ionic strength over the range 0.01 to 1 M, the presence of organic solvents including acetonitrile, methanol, ethanol, isopropanol, butanol, *N,N*-dimethylformamide and dimethylsulfoxide, the presence of detergents, and the presence of chaotropic agents (protein denaturants) including urea, guanidine hydrochloride, guanidine thiocyanate and formamide, which are incompatible with prior art indirect labeling systems wherein the structure of a biological macromolecule must be maintained to preserve requisite binding properties. Once formed, the bioconjugates are stable upon removal of water, and can be lyophilized for storage.

The stability of the bioconjugate at a given pH is determined to some extent by substituent group R_3 . Phenylboronic acid complexes of General Formula VI, wherein group R_3 is H, are stable in buffered aqueous solutions over the approximate pH range 3.5 to 10.5. Phenylboronic acid complexes of General Formula VI, wherein group R_3 is CH_3 , are stable in buffered aqueous solutions over the approximate pH range 4.5 to 10.5. Phenylboronic acid complexes of General Formula VI, wherein group R_3 includes an electronegative moiety, are stable in buffered aqueous solutions over the approximate pH range less than 3.5 to 10.5.


The stability of the phenylboronic acid complex toward acid catalyzed hydrolysis is related to the pK_a of the hydroxamic acid participating in the

-40-

complex. The lower the pK_a of the hydroxamic acid moiety the more stable the complex. Consequently, phenylboronic acid complexes of General Formula VI wherein group R_3 includes an electronegative moiety exhibit greater stability toward acid catalyzed hydrolysis than do those in which R_3 is either H or CH_3 .

The following examples present a detailed description of the synthesis of reagents of General Formula I and General Formula III.

Example I
Preparation of Ethyl 4-Aminomethylsalicylate
Hydrochloride



Ethyl 4-Methylsalicylate.

4-Methylsalicylic acid (20.0 g, 131 mmoles) was dissolved in ethanol (300 mL) and concentrated sulfuric acid (2.0 mL) was added. The mixture was refluxed for 40 hours. The volume of the reaction mixture was reduced to 100 mL, transferred to a separatory funnel, and diluted with chloroform (250 mL) and water (200 mL). Solid sodium bicarbonate was added in small portions until the pH of the aqueous layer was about 8 (pH test paper). The mixture in the funnel was shaken well and the layers separated. The organic layer was washed first with water (150 mL) and then saturated aqueous sodium chloride (150 mL). Finally, the organic solution was dried over anhydrous magnesium sulfate, filtered, and the solvent evaporated to afford 14.0 g (59% yield) of liquid ethyl 4-methylsalicylate.

-41-

^1H NMR (300 MHz, CHCl_3 - d) δ 1.40 (triplet, $J = 7$ Hz, 3H, CH_2CH_3), 2.33 (singlet, 3H, ArCH_3), 4.38 (quartet, $J = 7$ Hz, 2H, CH_2CH_3), 6.67 (doublet, $J = 8$ Hz, 1H, ArH), 6.78 (singlet, 1H, ArH), 7.72 (doublet, $J = 8$ Hz, 1H, ArH), 10.81 (singlet, 1H, OH). ^{13}C NMR (75 MHz, CHCl_3 - d) δ 14.0, 21.7, 61.1, 110.1, 117.7, 120.4, 129.7, 146.9, 161.7, 170.3.

Ethyl 4-Bromomethylsalicylic Acid.

Ethyl 4-methylsalicylate (13.1 g, 72.5 mmol) was dissolved in carbon tetrachloride (150 mL) and *N*-bromosuccinimide (13.1 g, 73.2 mmol) and benzoyl peroxide (0.2 g, 0.8 mmol) were added. The mixture was refluxed for 3.5 hours and then allowed to cool to room temperature. The solids were removed by filtration and the filtrate was evaporated to dryness. The crude solid product was crystallized from hexane (100 mL) to afford 5.0 g (27% yield) of ethyl 4-bromomethylsalicylic acid (m.p. 64-66°C).

^1H NMR (300 MHz, CHCl_3 - d) δ 1.41 (triplet, $J = 7$ Hz, 3H, CH_2CH_3), 4.40 (quartet, $J = 7$ Hz, 2H, CH_2CH_3), 4.40 (singlet, 2H, CH_2Br), 6.90 (doublet, $J = 8$ Hz, 1H, ArH), 6.99 (singlet, 1H, ArH), 7.82 (doublet, $J = 8$ Hz, 1H, ArH), 10.88 (singlet, 1H, OH). ^{13}C NMR (75 MHz, CHCl_3 - d) δ 14.0, 31.9, 61.5, 112.4, 117.9, 119.8, 130.5, 145.5, 161.8, 169.9.

Ethyl 4-Aminomethylsalicylate Hydrochloride.

Ethyl 4-bromomethylsalicylate (4.8 g, 18.6 mmol) was dissolved in dry *N,N*-dimethylformamide (50 mL) and sodium azide (1.2 g, 18.9 mmol) was added. The suspension was stirred at room temperature for 2 hours. The reaction mixture was then diluted with dichloromethane (150 mL) and extracted with 1 N aqueous hydrochloric acid (100 mL), water (100 mL), and saturated aqueous sodium chloride (50 mL). Finally, the solution was then dried over anhydrous

-42-


magnesium sulfate, filtered, and evaporated to dryness to give ethyl 4-azidomethylsalicylate as an oil.

Palladium on carbon (0.5 g; 10% [w/w]) was added to a 1L hydrogenation flask under a nitrogen atmosphere. The crude ethyl 4-azidomethylsalicylate was dissolved in ethanol (200 mL) and transferred to the hydrogenation flask. Concentrated aqueous hydrochloric acid (2 mL) was then added, and the flask was affixed to the Parr hydrogenator. The reaction mixture was shaken under 35 psi of hydrogen for 4 hours at room temperature. The mixture was then filtered through Celite to remove the catalyst, and the filtrate was evaporated to dryness to afford an off-white solid. Finally, this solid was crystallized from EtOH to afford 3.1 g (71 % yield) of ethyl 4-aminomethylsalicylate hydrochloride (m.p. 240 - 241 °C).

^1H NMR (300 MHz, DMSO- d_6) δ 1.30 (triplet, $J = 7$ Hz, 3H, CH_2CH_3), 3.99 (singlet, 2H, CH_2NH_3), 4.33 (quartet, $J = 7$ Hz, 2H, CH_2CH_3), 7.06 (doublet, $J = 8$ Hz, 1H, ArH), 7.15 (singlet, 1H, ArH), 7.77 (doublet, $J = 8$ Hz, 1H, ArH), 8.71 (broad singlet, 3H, NH_3), 10.62 (broad singlet, 1H, OH). ^{13}C NMR (75 MHz, DMSO- d_6) δ 14.0, 38.7, 41.6, 61.5, 112.9, 117.7, 119.8, 130.3, 142.2, 160.2, 168.9.

Example II

Preparation of a Reagent of General Formula I Ethyl *N*-Iodoacetyl-4-aminomethylsalicylate



Ethyl N-Chloroacetyl-4-aminomethylsalicylate.

Ethyl 4-aminomethylsalicylate hydrochloride (0.50 g, 2.17 mmoles) was suspended in dry *N,N*-dimethylformamide (25 mL) and *N,N*-diisopropylethylamine (0.38 mL, 2.18 mmoles) was added. Once the amine salt dissolved, chloroacetic anhydride (0.39 g, 2.25 mmoles) was added and the reaction mixture was stirred at room temperature for 4.5 hours. The reaction mixture was then diluted with ethyl acetate (100 mL), and this solution was extracted with 1 N aqueous hydrochloric acid (100 mL), water (50 mL), and saturated aqueous sodium chloride (50 mL). The ethyl acetate solution was dried over anhydrous magnesium sulfate, filtered and evaporated to dryness to afford a white solid. Finally, this solid was crystallized from ethyl acetate:hexanes (8:2, 10 mL) to afford 0.13 g (22% yield) of ethyl *N*-chloroacetyl-4-aminomethylsalicylate (m.p. 120-121°C).

¹H NMR (300 MHz, DMSO-*d*₆) δ 1.31, (triplet, *J* = 7 Hz, 3H, CH₂CH₃), 4.14 (singlet, 2H, ClCH₂), 4.29 (doublet, *J* = 6 Hz, 2H, NHCH₂), 4.34 (quartet, *J* = 7 Hz, 2H, CH₂CH₃), 6.82 (doublet, *J* = 8 Hz, 1H, ArH), 6.84 (singlet, 1H, ArH), 7.73 (doublet, *J* = 8 Hz, 1H, ArH), 8.78 (triplet, *J* = 6 Hz, 1H, NH), 10.58 (singlet, 1H, OH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 13.9, 42.1, 42.6, 61.3, 111.6, 115.6, 118.4, 130.1, 147.6, 160.6, 166.5, 169.1.

Ethyl N-Iodoacetyl-4-aminomethylsalicylate.


Ethyl *N*-chloroacetyl-4-aminomethylsalicylate (0.11 g, 0.40 mmoles) was dissolved in acetone (5 mL) and sodium iodide (0.06 g, 0.40 mmoles) was added. The solution was refluxed for 2.5 hours, then cooled to room temperature and filtered. The filtrate was evaporated to dryness to afford a white solid of ethyl *N*-iodoacetyl-4-aminomethylsalicylate (0.19 g, 100% yield) (m.p. 105 - 108°C).

-44-

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 1.31, (triplet, $J = 7$ Hz, 3H, CH_2CH_3), 3.68 (singlet, 2H, ICH_2), 4.26 (doublet, $J = 6$ Hz, 2H, NHCH_2), 4.33 (quartet, $J = 7$ Hz, 2H, CH_2CH_3), 6.82 (doublet, $J = 8$ Hz, 1H, ArH), 6.84 (singlet, 1H, ArH), 7.72 (doublet, $J = 8$ Hz, 1H, ArH), 8.79 (triplet, $J = 6$ Hz, 1H, NH), 10.58 (singlet, 1H, OH). ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 0.41, 13.9, 42.0, 61.3, 111.4, 115.5, 118.3, 130.1, 147.9, 160.6, 168.3, 169.1.

Example III

Preparation of a Reagent of General Formula I Ethyl (6-Aminohexanoyl)aminomethylsalicylate Trifluoroacetate



Ethyl (N-tert-Butoxycarbonyl-6-aminohexanoyl)aminomethylsalicylate.

Ethyl 4-aminomethylsalicylate hydrochloride (0.52 g, 1.28 mmoles) was suspended in anhydrous *N,N*-dimethylformamide (25 mL), and *N,N*-diisopropylethylamine (0.79 mL, 4.53 mmoles) was added, followed by *N*-tert-butoxycarbonyl-6-aminohexanoic acid succinimidyl ester (0.74 g, 2.26 mmoles). The mixture was stirred under dry nitrogen for 18 hours, during which time all solids dissolved. The reaction mixture was diluted with ethyl acetate (100 mL) and extracted with 1 N aqueous hydrochloric acid (100 mL). The layers were separated, and the ethyl acetate solution was washed with water (100 mL) and saturated aqueous sodium chloride (500 mL). The ethyl acetate solution was dried over anhydrous

-45-

magnesium sulfate, filtered, and evaporated to afford an amorphous off-white solid. Finally, the solid was crystallized from ethyl acetate, filtered, and dried *in vacuo* to afford 0.67 g (73% yield) of ethyl (*N*-*tert*-butoxycarbonyl-6-aminohexanoyl)aminomethylsalicylate (m.p. 120-121°C, open capillary, uncorrected).

¹H NMR (300 MHz, DMSO-*d*₆) δ 1.19 (multiplet, 2H, NHCH₂CH₂CH₂CH₂CH₂CO), 1.34 (multiplet, 5H, CH₂CH₂CO and CH₂CH₃), 1.34 (singlet, 9 H, C(CH₃)₃), 1.49 (multiplet, 2 H, NHCH₂CH₂), 2.12 (triplet, *J* = 7 Hz, 2 H, CH₂CH₂CO), 2.87 (quartet, *J* = 6 Hz, 2 H, NHCH₂CH₂), 4.23 (doublet, *J* = 6 Hz, 2 H, CH₂Ar), 4.32 (quartet, *J* = 7 Hz, CH₂CH₃), 6.74 (triplet, *J* = 6 Hz, 1 H, CONHCH₂CH₂), 6.80 (doublet, *J* = 8 Hz, 1 H, ArH), 6.81 (singlet, 1 H, ArH), 7.71 (doublet, *J* = 8 Hz, 1 H, ArH), 8.34 (triplet, *J* = 6 Hz, 1 H, CONHCH₂Ar), 10.58 (singlet, 1 H, OH).
¹³C NMR (75 MHz, DMSO-*d*₆) δ 14.0, 25.1, 26.3, 28.3, 29.7, 36.3, 40.2, 42.9, 61.4, 79.0, 111.6, 116.0, 118.3, 130.4, 146.9, 156.2, 161.9, 170.1, 173.0.

Ethyl (6-Aminohexanoyl)aminomethylsalicylate Trifluoroacetate.

Ethyl (*N*-*tert*-butoxycarbonyl-6-aminohexanoyl)aminomethylsalicylate (0.58 g, 1.41 mmoles) was dissolved in dichloromethane (5 mL) and the solution was cooled in an ice/water bath. Trifluoroacetic acid (5 mL) was added, and the reaction was allowed to warm to room temperature. After 2 hours, the reaction mixture was evaporated to dryness to give the product as an oil, which was dried *in vacuo* over potassium hydroxide pellets to afford 0.59 g (99% yield) of ethyl (6-aminohexanoyl)aminomethylsalicylate trifluoroacetate.

¹H NMR (300 MHz, DMSO-*d*₆) δ 1.28 (multiplet, 5H, H₃CH₂CH₂CH₂CH₂CH₂CO and CH₂CH₃), 1.53 (multiplet, 4H, NH₃CH₂CH₂CH₂CH₂CH₂CO), 2.15 (triplet, *J* = 8 Hz, 2 H,


-46-

CH₂CH₂CO), 2.71 (multiplet, 2 H, NH₃CH₂CH₂), 4.21 (doublet, $J = 6$ Hz, 2 H, CH₂Ar), 4.30 (quartet, $J = 8$ Hz, 2H, CH₂CH₃), 6.79 (doublet, $J = 8$ Hz, 1 H, ArH), 6.81 (singlet, 1 H, ArH), 7.68 (doublet, $J = 8$ Hz, 1 H, ArH), 8.18 (broad singlet, 3 H, NH₃), 8.60 (triplet, $J = 6$ Hz, 1 H, CONHCH₂Ar), 10.58 (broad singlet, 1 H, OH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 14.0, 24.8, 25.6, 26.7, 35.1, 38.6, 41.7, 61.3, 111.5, 115.5, 118.3, 130.1, 148.6, 160.6, 169.2, 172.6.

Example IV

Preparation of a Reagent of General Formula I

Methyl 4-Succinylaminomethylsalicylate Succinimidyl Ester



Methyl 4-Methylsalicylate.

4-Methylsalicylic acid (100 g, 658 mmoles) was dissolved in anhydrous methanol (500 mL) and concentrated sulfuric acid (25 mL) was added carefully. The solution was refluxed for 18 hours, then cooled to room temperature. The reaction mixture was concentrated to about 150 mL, and ethyl acetate (250 mL) was added. The ethyl acetate solution was washed twice with saturated aqueous sodium bicarbonate (250 mL portions) and then with saturated aqueous sodium chloride (100 mL). The ethyl acetate solution was dried over anhydrous sodium sulfate, filtered, and evaporated to a clear, reddish-brown liquid. This crude product was vacuum

-47-

distilled (oil pump) to afford a clear, viscous liquid that solidified on standing to afford 98.1 g (90% yield) of methyl 4-methylsalicylate.

^1H NMR (300 MHz, CHCl_3 -d) δ 2.32 (singlet, 3H, ArCH_3), 3.91 (singlet, 3H, OCH_3), 6.67 (doublet, $J = 8$ Hz, 1H, ArH), 6.78 (singlet, 1H, ArH), 7.69 (doublet, $J = 8$ Hz, 1H, ArH), 10.71 (singlet, 1H, OH). ^{13}C NMR (75 MHz, CHCl_3 -d) δ 21.8, 52.1, 110.0, 117.9, 120.6, 129.9, 147.3, 161.9, 170.9.

Methyl 4-Bromomethylsalicylate.

Methyl 4-methylsalicylate (98.1 g, 590 mmoles) was dissolved in carbon tetrachloride (600 mL), and *N*-bromosuccinimide (105.0 g, 590 mmoles) and benzoyl peroxide (0.7 g, 3 mmoles) were added. The mixture was refluxed under nitrogen. After 2 hours, an additional portion (0.7 g) of *N*-bromosuccinimide was added. Reflux was continued for 16 hours. The reaction mixture was cooled to room temperature and the solid removed by filtration. The yellow filtrate was evaporated to dryness to afford a thick yellow syrup that solidified on standing. Hexanes (500 mL) was added to the solid, and the mixture was boiled until almost all solids dissolved. The hot hexanes solution was filtered and concentrated until a solid just began to precipitate. The mixture was heated to dissolve the solid, and the solution was allowed to cool slowly to room temperature. Pale yellow crystals formed slowly. The mixture was then chilled in ice for 2 hours to complete crystallization. Finally, the solid was filtered, washed with cold hexanes (100 mL), and dried *in vacuo* to afford 83.5 g (58% yield) of methyl 4-bromomethylsalicylate (m.p. 73-75° C, open capillary, uncorrected).

^1H NMR (300 MHz, CHCl_3 -d) δ 3.95 (singlet, 3H, OCH_3), 4.40 (singlet, 2H, CH_2), 6.90 (doublet, $J = 8$ Hz, 1H, ArH), 7.00 (singlet, 1H, ArH), 7.80 (doublet, $J = 8$ Hz, 1H, ArH), 10.78

-48-

(singlet, 1H, OH). ^{13}C NMR (75 MHz, $\text{CHCl}_3\text{-}d$) δ 32.1, 52.5, 112.4, 118.2, 120.1, 130.7, 145.8, 162.0, 170.5.

Methyl 4-Azidomethylsalicylate.

Methyl 4-bromomethyl salicylate (83.5 g, 341 mmoles) was dissolved in dry *N,N*-dimethyl-formamide (150 mL) and sodium azide 925.0 g, 380 mmoles) was added. The yellow suspension was stirred at room temperature, and the solids rapidly dissolved. The solution turned brown, and a precipitate of sodium bromide formed. The reaction mixture was stirred 16 hours, then filtered. The filtrate was evaporated to a brown oil, which was dissolved in a mixture of hexanes and ethyl acetate (1:1 [v/v], 100 mL). Silica gel (25 g, flash chromatography grade) was added to the brown solution, and the mixture was swirled well. The silica was removed by filtration on a glass frit, and washed three times with hexanes:ethyl acetate (1:1 [v/v], 50 mL portions). The silica gel was dried on the frit, and the combined filtrates were evaporated to dryness to afford a dark yellow liquid. The crude product (which was utilized for the following reaction) was found to contain some residual *N,N*-dimethylformamide.

^1H NMR (300 MHz, $\text{CHCl}_3\text{-}d$) δ 3.94 (singlet, 3H, OCH_3), 4.32 (singlet, 2H, CH_2), 6.82 (doublet, $J = 8$ Hz, 1H, ArH), 6.93 (singlet, 1H, ArH), 7.83 (doublet, $J = 8$ Hz, 1H, ArH), 10.80 (singlet, 1H, OH). ^{13}C NMR (75 MHz, $\text{CHCl}_3\text{-}d$) δ 52.5, 54.3, 112.3, 117.0, 118.7, 130.8, 143.9, 162.1, 170.5.

Methyl 4-Aminomethylsalicylate Hydrochloride.

Crude methyl 4-azidomethylsalicylate was dissolved in methanol (750 mL) in a 2L Parr hydrogenation flask. Palladium on carbon catalyst (10% [w/w], 3.8 g) in water (25 mL) was added, followed by concentrated hydrochloric acid (35 mL). The flask was affixed to a Parr hydrogenator, and

-49-

the mixture was shaken at room temperature under 40 psi of hydrogen for 16 hours. The reaction mixture was then filtered through a 0.45 mm nylon filter. The retained solid was then washed with methanol (150 mL), water (100 mL), and methanol again (150 mL). The combined filtrates were evaporated to dryness to afford a tan solid. This solid was dissolved in hot denatured ethanol (150 mL) and the solution was allowed to cool to room temperature. White crystals formed quickly. Finally, the mixture was then chilled for 16-18 hours at 4° C to complete crystallization. The solid was filtered, washed with a little cold ethanol (50 mL) and then diethyl ether (150 mL), and dried *in vacuo* over potassium hydroxide pellets to afford 51.5 g (65% yield based on methyl 4-bromomethylsalicylate) of methyl 4-aminomethylsalicylate hydrochloride (m.p. 225 - 227° C, open capillary, uncorrected).

¹H (300 MHz, DMSO-*d*₆) δ 3.87 (singlet, 3H, OCH₃), 4.00 (singlet, 2H, CH₂), 7.06 (doublet, *J* = 8 Hz, 1H, ArH), 7.13 (singlet, 1H, ArH), 7.77 (doublet, *J* = 8 Hz, 1H, ArH), 8.59 (broad singlet, 3H, NH₃Cl), 10.55 (singlet, 1H, OH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 41.6, 52.6, 113.0, 117.7, 119.8, 130.4, 142.1, 160.0, 169.1

Methyl 4-Succinylaminomethylsalicylate.

Methyl 4-aminomethylsalicylate hydrochloride (3.00 g, 13.8 mmoles) was suspended in dry pyridine (25 mL), and *N,N*-diisopropylethylamine (2.7 mL, 15.5 mmoles) was added. The suspension was stirred in an ice/water bath for 15 minutes, and then succinic anhydride (1.36 g, 13.6 mmoles) was added. The mixture was allowed to warm to room temperature, during which time the starting solids dissolved. After stirring for 2 hours, the mixture was evaporated to dryness, and the resulting solid was partitioned between ethyl acetate (300 mL) and 1 M aqueous hydrochloric acid (100 mL). The layers were separated, and the ethyl acetate

-50-

solution was washed with 1 M hydrochloric acid (100 mL) and saturated aqueous sodium chloride (100 mL). The solution was dried over anhydrous sodium sulfate, filtered, and evaporated to dryness to yield a white solid. Finally, the solid was crystallized from ethyl acetate/hexanes, filtered, and dried *in vacuo* to afford 3.02 g (87% yield) of methyl 4-succinylaminomethylsalicylate (m.p. 161-163° C).

¹H NMR (300 MHz, DMSO-*d*₆) δ 2.43 (triplet, *J* = 6 Hz, 2H, CH₂CONH), 2.49 (triplet, *J* = 6 Hz, 2H, HO₂CCH₂), 3.87 (singlet, 3H, OCH₃), 4.27 (doublet, *J* = 6 Hz, 2H, ArCH₂NH), 6.83 (doublet, *J* = 8 Hz, 1H, ArH), 6.86 (singlet, 1H, ArH), 7.71 (doublet, *J* = 8 Hz, 1H, ArH), 8.43 (triplet, *J* = 6 Hz, 1H, NH), 10.54 (singlet, 1H, OH), 12.12 (singlet, 1H, CO₂H). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 29.1, 30.1, 41.9, 52.5, 111.3, 115.6, 118.4, 130.1, 148.7, 160.7, 169.7, 171.7, 174.2.

Methyl 4-Succinylaminomethylsalicylate Succinimidyl Ester.

Methyl 4-succinylaminomethylsalicylate (2.60 g, 10.2 mmoles) was dissolved in dry tetrahydrofuran (100 mL), and *N*-hydroxysuccinimide (1.29 g, 11.2 mmoles) and 1,3-dicyclohexylcarbodiimide (2.10 g, 10.2 mmoles) were added. The mixture was stirred at room temperature under dry nitrogen, and the solids rapidly dissolved. After about 20 minutes, a white precipitate formed. The reaction mixture was stirred 16-18 hours, then chilled several hours at -20° C. The mixture was filtered cold, and the solid washed with a little tetrahydrofuran (25 mL). The combined filtrates were evaporated to dryness, and the residue was crystallized from ethyl acetate/hexanes, filtered, and dried *in vacuo* to afford 2.39 g (62% yield) of methyl 4-succinylaminomethylsalicylate succinimidyl ester (m.p. 133-135° C).


¹H NMR (300 MHz, DMSO-*d*₆) δ 2.56 (triplet, *J* = 7 Hz, 2H, CH₂CONH), 2.80 (singlet, 4H, COCH₂CH₂CO), 2.91 (triplet, *J* = 7 Hz, 2H, NO₂CCH₂), 3.87 (singlet, 3H, OCH₃), 4.27 (doublet, *J* = 6

-51-

Hz, 2H, ArCH₂NH), 6.83 (doublet, $J = 9$ Hz, 1H, ArH), 6.84 (singlet, 1H, ArH), 7.71 (doublet, $J = 9$ Hz, 1H, ArH), 8.51 (triplet, $J = 6$ Hz, 1H, NH), 10.50 (singlet, 1H, OH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 25.4, 25.9, 29.2, 41.8, 52.4, 111.4, 115.6, 118.4, 130.2, 148.2, 160.4, 168.9, 169.4, 170.2, 170.4.

Example V

Preparation of a Reagent of General Formula I Methyl (6-Aminohexanoyl)aminomethylsalicylate Hydrochloride



Methyl (N-tert-Butoxycarbonyl-6-aminohexanoyl)aminomethylsalicylate.

Methyl 4-aminomethylsalicylate hydrochloride (2.25 g, 10.3 mmoles) was suspended in anhydrous *N,N*-dimethylformamide (30 mL), and *N,N*-diisopropylethylamine (3.6 mL, 20.7 mmoles) was added, followed by *N*-tert-butoxycarbonyl-6-aminohexanoic acid succinimidyl ester (3.38 g, 10.3 mmoles). The mixture was stirred under dry nitrogen for 18 hours, during which time all solids dissolved. The solvent was then evaporated to leave a light brown syrup, which was partitioned between ethyl acetate (100 mL) and 1 M aqueous hydrochloric acid (100 mL). The layers were separated, and the ethyl acetate solution was washed with saturated aqueous sodium bicarbonate (100 mL) and saturated aqueous sodium chloride (100 mL). The ethyl acetate solution was dried over anhydrous sodium sulfate, filtered, and

-52-

evaporated to an amorphous off-white solid. The solid was crystallized from ethyl acetate/hexanes, filtered, and dried *in vacuo* to afford 3.25 g (80% yield) of methyl (*N*-*tert*-butoxycarbonyl-6-aminohexanoyl)-aminomethylsalicylate (m.p. 120-121° C, open capillary, uncorrected).

¹H NMR (300 MHz, DMSO-*d*₆) δ 1.22 (multiplet, 2H, NHCH₂CH₂CH₂CH₂CH₂CO), 1.36 (multiplet, 2H, CH₂CH₂CO), 1.36 (singlet, 9 H, C(CH₃)₃), 1.51 (multiplet, 2 H, NHCH₂CH₂), 2.13 (triplet, *J* = 7 Hz, 2 H, CH₂CH₂CO), 2.87 (quartet, *J* = 6 Hz, 2 H, NHCH₂CH₂), 3.87 (singlet, 3 H, OCH₃), 4.24 (doublet, *J* = 6 Hz, 2 H, CH₂Ar), 6.75 (triplet, *J* = 6 Hz, 1 H, CONHCH₂CH₂), 6.80 (doublet, *J* = 8 Hz, 1 H, ArH), 6.82 (singlet, 1 H, ArH), 7.72 (doublet, *J* = 8 Hz, 1 H, ArH), 8.35 (triplet, *J* = 6 Hz, 1 H, CONHCH₂Ar), 10.53 (singlet, 1 H, OH). ¹³C NMR (75 MHz, CHCl₃-*d*) δ 25.0, 26.0, 28.2, 29.3, 35.3, 41.7, 52.4, 77.4, 111.2, 115.5, 118.3, 130.1, 148.7, 155.8, 160.5, 169.5, 172.6.

Methyl (6-Aminohexanoyl)aminomethylsalicylate Hydrochloride.


Methyl (*N*-*tert*-butoxycarbonyl-6-aminohexanoyl)aminomethylsalicylate (3.00 g, 7.60 mmoles) was dissolved in ethyl acetate (100 mL), and dry hydrogen chloride was bubbled slowly through the solution. The reaction mixture warmed as the gas dissolved. After 5 minutes, the gas was shut off, and the solution was stirred at room temperature. A white precipitate formed in the solution. After 30 minutes, the reaction mixture was chilled in ice for 2 hours, then the solid was filtered, washed with diethyl ether (50 mL) and dried *in vacuo* over potassium hydroxide pellets to afford 2.50 g (99% yield) of methyl (6-aminohexanoyl)aminomethylsalicylate hydrochloride (decomposes with effervescence at 158-160° C, open capillary, uncorrected).

-53-

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 1.28 (multiplet, 2H, $\text{H}_3\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CO}$), 1.54 (multiplet, 4H, $\text{NH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CO}$), 2.16 (triplet, $J = 8$ Hz, 2 H, $\text{CH}_2\text{CH}_2\text{CO}$), 2.71 (multiplet, 2 H, $\text{NH}_3\text{CH}_2\text{CH}_2$), 3.85 (singlet, 3 H, OCH_3), 4.22 (doublet, $J = 6$ Hz, 2 H, CH_2Ar), 6.80 (doublet, $J = 8$ Hz, 1 H, ArH), 6.83 (singlet, 1 H, ArH), 7.70 (doublet, $J = 8$ Hz, 1 H, ArH), 8.10 (broad singlet, 3 H, NH_3), 8.54 (triplet, $J = 6$ Hz, 1 H, CONHCH_2Ar), 10.35 (broad singlet, 1 H, OH). ^{13}C NMR (75 MHz, CHCl_3-d) δ 24.8, 25.6, 26.7, 35.1, 41.7, 111.5, 115.6, 118.4, 130.2, 148.6, 160.4, 169.4.

Example VI

Preparation of a Reagent of General Formula I Cyanomethyl 4-Glutarylaminomethylsalicylate Succinimidyl Ester



Methyl N-tert-Butoxycarbonylaminomethylsalicylate.

Methyl 4-aminomethylsalicylate hydrochloride (10.9 g, 50 mmoles) was suspended in anhydrous methanol (200 mL) and di-*tert*-butyldicarbonate (10.9 g, 50 mmoles) and triethylamine (7.0 mL, 50 mmoles) were added. The solid rapidly dissolved with the slow evolution of gas. The reaction mixture was stirred at room temperature for 18 hours under dry nitrogen, then evaporated to dryness to afford a white amorphous mass. This mass was partitioned between ethyl acetate (200 mL) and water (100 mL). The layers were

-54-

separated, and the ethyl acetate solution was dried over anhydrous sodium sulfate. The solution was filtered and evaporated to a white solid. This solid was crystallized from ethyl acetate/hexanes, filtered, and dried *in vacuo* to afford 13.7 g (97% yield) of methyl *N*-*tert*-butoxycarbonylaminomethyl-salicylate (m.p. 95-96° C, open capillary, uncorrected).

¹H NMR (300 MHz, CHCl₃-*d*) δ 1.42 (singlet, 9H, C(CH₃)₃), 3.90 (singlet, 3H, OCH₃), 4.26 (doublet, *J* = 6 Hz, 2 H, CH₂Ar), 4.99 (triplet, *J* = 6 Hz, 1H, NH), 6.75 (doublet, *J* = 8 Hz, 1 H, ArH), 6.84 (singlet, 1 H, ArH), 7.73 (doublet, *J* = 8 Hz, 1 H, ArH), 10.72 (singlet, 1 H, OH). ¹³C NMR (75 MHz, CHCl₃-*d*) δ 28.5, 44.4, 52.4, 80.0, 111.5, 115.9, 118.2, 130.5, 150.0, 156.3, 162.1, 170.8.

N-*tert*-Butoxycarbonylaminomethylsalicylic acid.

Methyl *N*-*tert*-butoxycarbonylaminomethylsalicylate (8.7 g, 30.9 mmoles) was dissolved in dry tetrahydrofuran (100 mL), and potassium trimethylsilanolate (4.4 g, 30.9 mmoles, 90% pure) was added. The yellow solution was refluxed for 24 hours, during which time a tan precipitate formed and the solvent turned light brown. The mixture was evaporated to dryness, and the solid was dissolved in cold water (100 mL). The brown solution was chilled in an ice bath, and saturated aqueous potassium hydrogen sulfate solution was used to titrate the stirred solution to pH 2-3. An off-white solid precipitated during the titration. The solid was filtered, washed with cold water, and dried *in vacuo* over potassium hydroxide to afford 6.7 g, (81% yield) of crude *N*-*tert*-butoxycarbonylaminomethylsalicylic acid (m.p. 141-144° C, decomposes with effervescence on melting, open capillary, uncorrected).

¹H NMR (300 MHz, CHCl₃-*d*) δ 1.47 (singlet, 9H, C(CH₃)₃), 4.33 (doublet, *J* = 6 Hz, 2 H, CH₂Ar), 5.07 (triplet, *J* = 6 Hz, 1H,

-55-

NH), 6.80 (doublet, $J = 8$ Hz, 1 H, ArH), 6.88 (singlet, 1 H, ArH), 7.81 (doublet, $J = 8$ Hz, 1 H, ArH), 10.72 (broad singlet, 2 H, OH and CO₂H). ¹³C NMR (75 MHz, CHCl₃-*d*) δ 28.5, 44.4, 80.5, 111.1, 115.9, 118.3, 131.5, 148.5, 156.6, 162.6, 174.1.

Cyanomethyl N-tert-Butoxycarbonylaminomethylsalicylate.

*N-tert-Butoxycarbonyl*amino-methylsalicylic acid (8.2 g, 30.6 mmoles) was suspended in chloroacetonitrile (25 mL), and triethylamine (4.3 mL, 30.6 mmoles) was added. The mixture was stirred under dry nitrogen at 50° C, and the solids dissolved. The solution was stirred 18 hours, and then cooled to room temperature. The solvent was evaporated, and the residue was partitioned between ethyl acetate (250 mL) and water (250 mL). The layers were separated, and the ethyl acetate layer was washed with saturated aqueous sodium bicarbonate (100 mL) and saturated aqueous sodium chloride (100 mL). The solution was dried over anhydrous sodium sulfate, filtered, and evaporated to dryness. The residual pale tan solid was dissolved in ethyl acetate (100 mL), and silica gel (10 g, flash chromatography grade) was added. The mixture was swirled well and allowed to sit for five minutes at room temperature. The silica was removed by filtration on a glass frit, and washed with ethyl acetate (100 mL). The filtrate was evaporated to dryness. The solid residue was crystallized from ethyl acetate/hexanes to afford 7.9 g (88% yield) of cyanomethyl *N-tert-butoxycarbonylaminomethylsalicylate* (m.p. 144-146° C, open capillary, uncorrected).

¹H NMR (300 MHz, CHCl₃-*d*) δ 1.45 (singlet, 9H, C(CH₃)₃), 4.30 (doublet, $J = 6$ Hz, 2 H, CH₂Ar), 5.00 (singlet, 2H, OCH₂CN), 5.05 (triplet, $J = 6$ Hz, 1H, NH), 6.83 (doublet, $J = 8$ Hz, 1 H, ArH), 6.91 (singlet, 1 H, ArH), 7.77 (doublet, $J = 8$ Hz, 1 H, ArH), 10.12 (singlet, 1 H, OH). ¹³C NMR (75 MHz, CHCl₃-*d*) δ

-56-

28.4, 44.3, 49.0, 80.1, 109.6, 114.2, 116.1, 118.7, 130.5, 149.7, 156.2, 162.5, 168.5.

Cyanomethyl Aminomethylsalicylate Hydrochloride.

Cyanomethyl *N*-*tert*-butoxycarbonylaminomethylsalicylate (7.7 g, 26.2 mmoles) was dissolved in tetrahydrofuran (150 mL), and dry hydrogen chloride was bubbled slowly through the solution. The reaction mixture warmed as the gas dissolved. After 5 minutes, the gas was shut off, and the solution was stirred at room temperature. A thick, creamy white precipitate formed in the solution. After 30 minutes, the reaction mixture was chilled in ice for 2 hours, then the solid was filtered, washed with diethyl ether (100 mL) and dried *in vacuo* over potassium hydroxide pellets to afford 5.8 g (91% yield) of cyanomethyl aminomethylsalicylate hydrochloride (m.p. darkens at 210° C, decomposes at 228° C, open capillary, uncorrected).

¹H NMR (300 MHz, DMSO-*d*₆) δ 4.00 (singlet, 2 H, CH₂Ar), 5.20 (singlet, 2H, OCH₂CN), 7.05 (doublet, *J* = 8 Hz, 1 H, ArH), 7.15 (singlet, 1 H, ArH), 7.75 (doublet, *J* = 8 Hz, 1 H, ArH), 8.62 (broad singlet, 3H, NH₃), 10.38 (singlet, 1 H, OH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 41.6, 49.7, 113.0, 116.1, 118.0, 119.7, 131.1, 142.3, 159.5, 166.1.

Cyanomethyl 4-Glutarylaminomethylsalicylate.

Cyanomethyl 4-aminomethylsalicylate hydrochloride (1.22 g, 5.0 mmoles) was suspended in dry dichloromethane (100 mL), and the suspension was stirred in an ice/water bath. A solution of glutaric anhydride (0.57 g, 5.0 mmoles) and triethylamine (0.7 mL, 5.0 mmoles) in dry dichloromethane (25 mL) was then added dropwise over 15 minutes. The mixture was allowed to warm to room temperature, and the reaction was stirred for 18 hours. The mixture was evaporated to dryness, and the resulting solid was triturated

-57-

under cold 0.1 M aqueous hydrochloric acid (50 mL). The solid was collected by filtration, washed with cold water, and dried *in vacuo* over potassium hydroxide pellets to afford 1.43 g (89% yield) of cyanomethyl 4-glutarylaminomethylsalicylate (m.p. 125-126° C).

¹H NMR (300 MHz, DMSO-*d*₆) δ 1.75 (quintet, *J* = 7 Hz, 2H, CH₂CH₂CH₂), 2.19 (triplet, *J* = 7 Hz, 2H, CH₂CONH), 2.22 (triplet, *J* = 7 Hz, 2H, HO₂CCH₂), 4.24 (doublet, *J* = 6 Hz, 2H, ArCH₂NH), 5.17 (singlet, 2H, OCH₂CN), 6.81 (doublet, *J* = 8 Hz, 1H, ArH), 6.85 (singlet, 1H, ArH), 7.70 (doublet, *J* = 8 Hz, 1H, ArH), 8.39 (triplet, *J* = 6 Hz, 1H, NH), 10.5 (very broad singlet, 1H, OH), 11.7 (very broad singlet, 1H, CO₂H). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 20.7, 33.1, 34.5, 41.8, 49.7, 111.2, 115.9, 116.2, 118.5, 130.9, 149.1, 160.2, 166.6, 172.3, 174.5.

Cyanomethyl 4-Glutarylaminomethyl salicylate Succinimidyl Ester.

Cyanomethyl 4-glutarylaminomethylsalicylate (1.00 g, 3.1 mmoles) was dissolved in dry tetrahydrofuran (50 mL), and *N*-hydroxysuccinimide (0.36 g, 3.1 mmoles) and 1,3-dicyclohexylcarbodiimide (0.64 g, 3.1 mmoles) were added. The mixture was stirred at room temperature under dry nitrogen, and the solids rapidly dissolved. After about 60 minutes, a white precipitate formed. The reaction mixture was stirred 24 hours, then chilled several hours at -20° C. The mixture was filtered cold, and the solid washed with a little tetrahydrofuran (10 mL). The combined filtrates were evaporated to dryness, and the residue was crystallized from ethyl acetate/hexanes, filtered, and dried *in vacuo* to afford 1.04 g (80% yield) of cyanomethyl 4-glutarylaminomethylsalicylate succinimidyl ester (m.p. 116-119° C).


¹H NMR (300 MHz, DMSO-*d*₆) δ 1.88 (quintet, *J* = 7 Hz, 2H, CH₂CH₂CH₂), 2.30 (triplet, *J* = 7 Hz, 2H, CH₂CONH), 2.71

-58-

(triplet, $J = 7$ Hz, 2H, NO_2CCH_2), 2.80 (singlet, 4H, $\text{COCH}_2\text{CH}_2\text{CO}$), 4.26 (doublet, $J = 6$ Hz, 2H, ArCH_2NH), 5.18 (singlet, 2H, OCH_2CN), 6.82 (doublet, $J = 8$ Hz, 1H, ArH), 6.86 (singlet, 1H, ArH), 7.71 (doublet, $J = 8$ Hz, 1H, ArH), 8.44 (triplet, $J = 6$ Hz, 1H, NH), 10.18 (broad singlet, 1H, OH). ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 20.4, 25.5, 29.7, 33.6, 41.8, 49.6, 111.2, 115.9, 116.2, 118.4, 130.9, 148.9, 160.1, 166.5, 169.0, 170.5, 171.7.

Example VII

Preparation of a Reagent of General Formula I Cyanomethyl 4-(6-maleimidohexanoyl)aminomethylsalicylate



Cyanomethyl 4-(6-maleimidohexanoyl)aminomethylsalicylate.

Cyanomethyl 4-aminomethylsalicylate hydrochloride (79 mg, 0.32 mmoles) and 6-maleimidocaproic acid *N*-hydroxysuccinimide ester (100 mg, 0.32 mmoles) were suspended in dry *N,N*-dimethylformamide (5.0 mL), and the suspension was stirred at room temperature. *N,N*-diisopropylethylamine (87 μL , 0.50 mmoles) was added. The solids rapidly dissolved to afford a clear, pale yellow solution. After 30 minutes, the mixture was evaporated to dryness, and the residue was partitioned between ethyl acetate (25 mL) and cold 0.1 M aqueous hydrochloric acid (25 mL). The layers were separated, and the ethyl acetate


-59-

solution washed with saturated aqueous sodium bicarbonate (25 mL) and saturated aqueous sodium chloride (25 mL). The ethyl acetate solution was dried over anhydrous sodium sulfate, filtered and evaporated to a white solid. The solid was crystallized from ethyl acetate/hexanes to afford 108 mg (84% yield) of cyanomethyl 4-(6-maleimidohexanoyl)aminomethyl salicylate (m.p. 141-144°C).

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 1.20 (multiplet, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$), 1.53 (multiplet, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$), 2.13 (triplet, $J = 7$ Hz, 2H, NCH_2CH_2), 3.38 (triplet, $J = 7$ Hz, 2H, $\text{CH}_2\text{CH}_2\text{CO}$), 4.26 (doublet, $J = 6$ Hz, 2H, ArCH_2NH), 5.02 (singlet, 2H, OCH_2CN), 6.68 (singlet, 2H, $\text{CH}=\text{CH}$), 6.76 (doublet, $J = 8$ Hz, 1H, ArH), 6.80 (singlet, 1H, ArH), 7.69 (doublet, $J = 8$ Hz, 1H, ArH), 7.99 (triplet, $J = 6$ Hz, 1H, NH), 10.04 (singlet, 1H, OH). ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 24.5, 25.7, 27.6, 35.2, 36.9, 41.9, 48.8, 109.1, 114.3, 115.6, 118.2, 129.8, 133.8, 149.2, 161.2, 167.6, 170.4, 172.7.

Example VIII

Preparation of a Reagent of General Formula I Cyanomethyl 4-(3-(2-Pyridyldithio)propionyl)aminomethylsalicylate



Cyanomethyl 4-(3-(2-pyridyldithio)propionyl)aminomethylsalicylate.

Cyanomethyl 4-aminomethylsalicylate hydrochloride (79 mg, 0.32 mmoles) and 3-(2-pyridyldithio)propionic acid *N*-


-60-

hydroxysuccinimide ester (100 mg, 0.32 mmoles) were suspended in dry *N,N*-dimethylformamide (5.0 mL), and the suspension was stirred at room temperature. *N,N*-Diisopropylethylamine (87 μ L, 0.50 mmoles) was added. The solids rapidly dissolved to give a clear, pale tan solution. After 30 minutes, the mixture was evaporated to dryness, and the residue was partitioned between ethyl acetate (25 mL) and cold 0.1 M aqueous hydrochloric acid (25 mL). The layers were separated and the ethyl acetate solution washed with saturated aqueous sodium bicarbonate (25 mL) and saturated aqueous sodium chloride (25 mL). The ethyl acetate solution was dried over anhydrous sodium sulfate, filtered and evaporated to a clear oil. Trituration under cold hexanes afforded 64 mg (48% yield) of a gum of cyanomethyl 4-(3-(2-pyridyldithio)propionyl)aminomethylsalicylate.

^1H NMR (300 MHz, DMSO- d_6) δ 2.60 (triplet, $J = 7$ Hz, 2H, SCH_2CH_2), 3.05 (triplet, $J = 7$ Hz, 2H, $\text{CH}_2\text{CH}_2\text{CO}$), 4.27 (doublet, $J = 6$ Hz, 2H, ArCH_2NH), 5.19 (singlet, 2H, OCH_2CN), 6.84 (doublet, $J = 8$ Hz, 1H, ArH), 6.88 (singlet, 1H, ArH), 7.23 (triplet, $J = 6$ Hz, 1H, ArH), 7.71 (doublet, $J = 8$ Hz, 1H, ArH), 7.74-7.82 (multiplet, 2H, ArH), 8.44 (doublet, $J = 6$ Hz, 1H, ArH), 8.56 (triplet, $J = 6$ Hz, 1H, NH), 10.13 (broad singlet, 1H, OH). ^{13}C NMR (75 MHz, DMSO- d_6) δ 34.0, 34.5, 41.8, 49.6, 54.9, 111.2, 115.9, 116.0, 118.4, 119.3, 121.3, 130.8, 137.9, 148.6, 149.7, 159.4, 160.0, 166.4, 170.3.

-61-

Example IX
Preparation of a Reagent of General Formula III
4-Glutarylaminomethylsalicylhydroxamic Acid
Hydrazide



Methyl 4-Glutarylaminomethylsalicylate N-tert-Butyloxycarbonylhydrazide.

Methyl 4-glutarylaminomethylsalicylate succinimidyl ester (2.57 g, 6.8 mmoles) [prepared as for the succinyl derivative above, substituting glutaric anhydride for succinic anhydride] was dissolved in dry tetrahydrofuran (100 mL), and *tert*-butylcarbazate (0.90 g, 6.8 mmoles) was added. The reaction was stirred at room temperature for 60 hours. The solution was then evaporated to dryness, and the residue dissolved in ethyl acetate (100 mL). The ethyl acetate solution was washed with saturated aqueous potassium bicarbonate (100 mL) and saturated aqueous sodium chloride (100 mL). It was then dried over anhydrous sodium sulfate, filtered, and evaporated to dryness. The residue was crystallized by dissolving it in ethyl acetate (50 mL) with gentle warming, adding hexanes (50 mL) to the warm solution, and chilling at -20° C. Once crystallization began, another portion of hexanes (50 mL) was added, and the mixture was chilled for 18 hours at -20° C. Finally, the solid was filtered, washed with hexanes:ethyl acetate (2:1 [v/v], 90 mL), and dried *in vacuo* to afford 2.51 g (90% yield) of methyl 4-glutarylaminomethyl-salicylate *N-tert-*

-62-

butyloxycarbonylhydrazide (m.p. 68-72° C, open capillary, uncorrected).

¹H NMR (300 MHz, DMSO-*d*₆) δ 1.37 (singlet, 9H, (CH₃)₃C), 1.74 (quintet, *J* = 7 Hz, 2H, CH₂CH₂CH₂), 2.07 (triplet, *J* = 7 Hz, 2H, CH₂CONHCH₂), 2.17 (triplet, *J* = 7 Hz, 2H, HNHNOCCCH₂), 3.86 (singlet, 3H, OCH₃), 4.23 (doublet, *J* = 6 Hz, 2H, ArCH₂NH), 6.80 (doublet, *J* = 8 Hz, 1H, ArH), 6.82 (singlet, 1H, ArH), 7.71 (doublet, *J* = 8 Hz, 1H, ArH), 8.37 (triplet, *J* = 6 Hz, 1H, NH), 8.65 (singlet, 1H, NHNHCOCH₂), 9.48 (singlet, 1H, OCONHNH), 10.50 (singlet, 1H, OH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 21.2, 28.0, 32.6, 34.6, 41.7, 52.4, 79.1, 111.4, 115.5, 118.3, 130.2, 148.5, 155.5, 160.4, 169.4, 171.7, 172.1.

4-Glutarylaminomethylsalicylhydroxamic Acid N-tert-Butyloxycarbonylhydrazide.

Methyl 4-glutarylaminomethylsalicylate *N-tert*-butyloxycarbonylhydrazide (2.00 g, 4.9 mmoles) was added to a cooled (ice/water bath) solution of hydroxylamine sulfate (0.82 g, 5.0 mmoles), sodium hydroxide (1.00 g, 25.0 mmoles) and sodium sulfite (0.20 g, mmoles) in water (25 mL). The suspension was stirred for 18 hours in the dark, allowing it to warm to room temperature. The solution was then filtered to remove some insoluble material, and the pale yellow solution was chilled in an ice/water bath. The cold reaction mixture was slowly titrated to pH 3-4 with cold sulfuric acid (25% [v/v] aqueous), during which time a gummy material precipitated. The mixture was then chilled several hours in ice, and the liquid was decanted. The remaining amorphous solid was dissolved in methanol (25 mL), filtered, and evaporated to a white foam. The foam was dried *in vacuo* to 1.86 g (94% yield) of crude 4-glutarylaminomethylsalicylhydroxamic acid *N-tert*-butyloxycarbonylhydrazide.

¹H NMR (300 MHz, DMSO-*d*₆) δ 1.37 (singlet, 9H, (CH₃)₃C), 1.74 (quintet, *J* = 7 Hz, 2H, CH₂CH₂CH₂), 2.10 (triplet, *J* = 7 Hz,

-63-

2H, CH₂CONHCH₂), 2.16 (triplet, $J = 7$ Hz, 2H, H₂NHNOCCCH₂), 4.20 (doublet, $J = 6$ Hz, 2H, ArCH₂NH), 6.71 (doublet, $J = 8$ Hz, 1H, ArH), 6.74 (singlet, 1H, ArH), 7.60 (doublet, $J = 8$ Hz, 1H, ArH), 8.32 (triplet, $J = 6$ Hz, 1H, NH), 8.65 (singlet, 1H, NHNHCOCH₂), 9.29 (broad singlet, 1H, NHOH), 9.48 (singlet, 1H, OCONHNH), 11.37 (broad singlet, 1H, ArOH), 12.32 (broad singlet, 1H, NHOH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 21.2, 28.0, 32.6, 34.6, 41.7, 79.1, 112.4, 115.6, 117.6, 127.1, 145.9, 155.6, 159.8, 166.5, 171.7, 172.1.

4-Glutarylaminomethylsalicylhydroxamic Acid Hydrazide Hydrochloride.


4-Glutarylaminomethylsalicylhydroxamic acid *N*-*tert*-butyloxycarbonylhydrazide (1.86 g, 4.6 mmol) was dissolved in dry tetrahydrofuran (100 mL), and dry hydrogen chloride was bubbled slowly through the solution. The reaction mixture warmed as the gas dissolved. After 5 minutes, the gas was shut off, and the solution was stirred at room temperature. A white precipitate formed in the solution. After 30 minutes, the reaction mixture was chilled in ice for 2 hours, then the solid was filtered, washed with diethyl ether (50 mL) and dried *in vacuo* over potassium hydroxide pellets to afford 1.51 g (95% yield) of 4-glutarylaminomethylsalicylhydroxamic acid hydrazide hydrochloride (m.p. shrinks at 65° C, decomposes with effervescence at 100° C, open capillary, uncorrected).

¹H NMR (300 MHz, DMSO-*d*₆) δ 1.72 (quintet, $J = 7$ Hz, 2H, CH₂CH₂CH₂), 2.17 (triplet, $J = 7$ Hz, 2H, CH₂CONHCH₂), 2.23 (triplet, $J = 7$ Hz, 2H, H₂NHNOCCCH₂), 4.17 (doublet, $J = 6$ Hz, 2H, ArCH₂NH), 6.69 (doublet, $J = 8$ Hz, 1H, ArH), 6.74 (singlet, 1H, ArH), 7.66 (doublet, $J = 8$ Hz, 1H, ArH), 8.49 (triplet, $J = 6$ Hz, 1H, NH), 10.49 (broad singlet, 4H, NH₃NHCOCH₂), 11.08 (broad singlet, 1H, ArOH), 11.43 (broad singlet, 2H, NHOH). ¹³C NMR

-64-

(75 MHz, DMSO- d_6) δ 21.0, 25.2, 32.3, 34.4, 41.8, 67.2, 112.6, 115.7, 117.7, 127.4, 145.9, 159.8, 166.5, 171.7, 172.1.

Example X
Preparation Of A Reagent Of General Formula III
4-Glutarylaminomethylsalicyl(O-methyl)hydroxamic
Acid Succinimidyl Ester



Methyl N-(Benzyloxycarbonyl)-4-aminomethylsalicylate.

Methyl 4-aminomethylsalicylate hydrochloride (5.04 g, 23.2 mmoles) was suspended in chloroform (80 mL) and *N,N*-diisopropylethylamine (4.10 mL, 23.5 mmoles) and *N*-(benzyloxycarbonyloxy)succinimide (6.48 g, 26.0 mmoles) were added. The reaction mixture was stirred at room temperature for 4 hours, during which time all solids dissolved. The reaction mixture was then extracted with 1 N aqueous hydrochloric acid (100 mL), water (75 mL), and saturated aqueous sodium chloride (50 mL). The chloroform solution was dried over anhydrous magnesium sulfate, filtered and evaporated to dryness to give a white solid. The product was crystallized from ethyl acetate:hexanes to afford 6.17 g (84% yield) of methyl *N*-(benzyloxy-carbonyl)-4-aminomethylsalicylate (m.p. 91-92° C, open capillary, uncorrected).

^1H NMR (300 MHz, DMSO- d_6) δ 3.86 (singlet, 3H, OCH₃), 4.21 (doublet, J = 6 Hz, 2H, NHCH₂), 5.05 (singlet, 2H, CH₂O),

-65-

6.82 (doublet, $J = 8$ Hz, 1H, ArH), 6.86 (singlet, 1H, ArH), 7.27 - 7.36 (multiplet, 5H, ArH), 7.72 (doublet, $J = 8$ Hz, 1H, ArH), 7.90 (triplet, $J = 6$ Hz, 1H, NH), 10.53 (singlet, 1H, OH). ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 43.5, 52.4, 65.6, 111.5, 115.4, 118.2, 127.9, 128.0, 128.5, 130.2, 137.3, 148.6, 156.7, 160.5, 169.5.

Methyl N-(Benzyloxycarbonyl)-4-aminomethyl-2-O-benzylsalicylate.

Methyl *N*-(benzyloxycarbonyl)-4-aminomethylsalicylate (6.06 g, 19.2 mmoles) was dissolved in acetone (150 mL), and benzyl bromide (2.60 mL, 21.9 mmoles) and anhydrous potassium carbonate (13.28 g, 96.1 mmoles) were added. The mixture was stirred and heated at reflux for 22 hours. The mixture was concentrated to remove most of the acetone, and ethyl acetate (100 mL) was added. Aqueous hydrochloric acid (1N, 200 mL) was added slowly, swirling frequently to dissolve the solid carbonate. The layers were separated, and the aqueous layer was extracted with ethyl acetate (50 mL). The ethyl acetate solutions were combined and washed with water (100 mL) and saturated aqueous sodium chloride (50 mL), dried over anhydrous magnesium sulfate, filtered, and concentrated to about 50 mL. This solution was heated to boiling, and hexanes (150 mL) were added. The solution was cooled in ice, and crystals slowly formed. The crystals were collected by filtration, washed with hexanes, and dried *in vacuo* to afford 6.97 g (89% yield) of methyl *N*-(benzyloxycarbonyl)-4-aminomethyl-2-*O*-benzylsalicylate (m.p. 106-107° C, open capillary, uncorrected).

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 3.78 (singlet, 3H, OCH_3), 4.27 (doublet, $J = 6$ Hz, 2H, NHCH_2), 5.07 (singlet, 2H, CH_2OCO), 5.15 (singlet, 2H, CH_2O), 6.94 (doublet, $J = 8$ Hz, 1H, ArH), 7.16 (singlet, 1H, ArH), 7.26 - 7.42 (multiplet, 8H, ArH), 7.49 - 7.51 (multiplet, 2H, ArH), 7.68 (doublet, $J = 8$ Hz, 1H, ArH), 7.90 (triplet, $J = 6$ Hz, 1H, NH). ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ

-66-

43.8, 51.8, 65.6, 69.7, 112.5, 112.6, 118.9, 119.0, 127.2, 127.9, 128.0, 128.5 (2 carbons), 131.3, 137.0, 137.3, 146.2, 156.7, 157.8, 166.1.

N-(Benzyloxycarbonyl)-4-aminomethyl-2-*O*-benzylsalicylic acid.

Methyl *N*-(benzyloxycarbonyl)-4-aminomethyl-2-*O*-benzylsalicylate (6.81 g, 16.8 mmoles) was dissolved in tetrahydrofuran (50 mL). A solution of lithium hydroxide monohydrate (0.78 g, 18.5 mmoles) in water (25 mL) was added, and the reaction mixture was stirred and heated at 75°C for 24 hours. The solution was cooled, and 1 N aqueous hydrochloric acid (50 mL) was added. The reaction mixture was extracted twice with ethyl acetate (150 mL then 50 mL). The combined ethyl acetate extracts were washed with water (75 mL) and saturated aqueous sodium chloride (50 mL), dried over anhydrous magnesium sulfate, filtered, and concentrated to 150 mL. The ethyl acetate solution was heated to boiling, and hexanes (150 mL) was added. The solution was cooled in ice, and crystals formed. The crystals were collected by filtration, washed with hexanes, and dried *in vacuo* to afford 5.92 g (90% yield) of *N*-(benzyloxy-carbonyl)-4-aminomethyl-2-*O*-benzylsalicylic acid (m.p. 139-140° C, open capillary, uncorrected).

¹H NMR (300 MHz, DMSO-*d*₆) δ 4.23 (doublet, *J* = 6 Hz, 2H, NHCH₂), 5.06 (singlet, 2H, CH₂OCO), 5.13 (singlet, 2H, CH₂O), 6.90 (doublet, *J* = 8 Hz, 1H, ArH), 7.11 (singlet, 1H, ArH), 7.28 - 7.40 (multiplet, 8H, ArH), 7.48 - 7.51 (multiplet, 2H, ArH), 7.64 (doublet, *J* = 8 Hz, 1H, ArH), 7.88 (triplet, *J* = 6 Hz, 1H, NH), 12.57 (singlet, 1H, COOH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 43.8, 65.6, 69.8, 112.5 (2 carbons), 118.9, 120.4, 127.4, 127.9, 128.0, 128.5 (2 carbons), 131.2, 137.1, 137.4, 145.6, 156.7, 157.5, 167.4.

-67-

N-(Benzyloxycarbonyl)-4-aminomethyl-2-*O*-benzylsalicyl(*O*-methyl)hydroxamic acid.

N-(Benzyloxycarbonyl)-4-aminomethyl-2-*O*-benzylsalicylic acid (3.07 g, 7.84 mmoles) was dissolved in anhydrous *N,N*-dimethylformamide (75 mL) under dry nitrogen. After cooling the solution to in an ice/water bath, triethylamine (2.20 mL, 15.8 mmoles) was added followed by isobutyl chloroformate (1.10 mL, 8.48 mmoles). The reaction was stirred in ice for 1.5 hours. Methoxylamine hydrochloride (0.67 g, 8.02 mmoles) was added and the reaction mixture was allowed to warm to room temperature. After 3 hours, the reaction was diluted with ethyl acetate (150 mL) and extracted with 1 N aqueous hydrochloric acid (100 mL), water (100 mL), and saturated aqueous sodium chloride (50 mL). The ethyl acetate solution was dried over anhydrous magnesium sulfate, filtered and concentrated to about 50 mL. This solution was boiled and hexanes (100 mL) were added. Upon cooling in an ice bath, crystals formed quickly. They were collected by filtration and washed with hexane to afford 2.87 g (87% yield) of *N*-(benzyloxy-carbonyl)-4-aminomethyl-2-*O*-benzylsalicyl(*O*-methyl)hydroxamic acid (m.p. 116-117° C, open capillary, uncorrected).

¹H NMR (300 MHz, DMSO-*d*₆) δ 3.63 (singlet, 3H, NHOCH₃), 4.20 (doublet, *J* = 6 Hz, 2H, NHCH₂), 5.04 (singlet, 2H, CH₂OCO), 5.13 (singlet, 2H, CH₂O), 6.90 (doublet, *J* = 8 Hz, 1H, ArH), 7.08 (singlet, 1H, ArH), 7.29 - 7.49 (multiplet, 11H, ArH), 7.88 (triplet, *J* = 6 Hz, 1H, NH), 11.13 (singlet, 1H, NHOCH₃). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 43.7, 63.2, 65.5, 69.8, 111.8, 119.2, 121.7, 127.6, 127.9, 128.0, 128.1, 128.5 (2 carbons), 129.8, 136.8, 137.3, 144.3, 155.9, 156.6, 163.3.

-68-

4-Aminomethylsalicyl(O-methyl)hydroxamic Acid Hydrochloride.

N-(Benzyloxycarbonyl)-4-aminomethyl-2-*O*-benzylsalicyl(*O*-methyl)hydroxamic acid (1.42 g, 3.38 mmoles) and palladium on carbon (0.10 g, 10% [w/w]) were placed in a 1L hydrogenation flask under dry nitrogen. Ethanol (150 mL) was added, followed by concentrated aqueous hydrochloric acid (0.30 mL). The flask was affixed to the Parr hydrogenator and shaken under 35 psi of hydrogen for 7 hours at room temperature. The reaction mixture was then filtered through Celite to remove the catalyst, and the filtrate was concentrated until a precipitate began to form. The mixture was cooled in ice, the solid collected by filtration, washed with hexanes, and dried *in vacuo* to afford 0.65 g (83% yield) of 4-aminomethyl-salicyl(*O*-methyl)-hydroxamic acid hydrochloride (m.p. > 250° C, open capillary, uncorrected).

¹H NMR (300 MHz, DMSO-*d*₆) δ 3.71 (singlet, 3H, NHOCH₃), 3.95 (singlet, 2H, NH₃CH₂), 7.00 (doublet, *J* = 8 Hz, 1H, ArH), 7.05 (singlet, 1H, ArH), 7.72 (doublet, *J* = 8 Hz, H, ArH), 8.53 (broad singlet, 3H, NH₃), 12.01 (broad singlet, 2H, NHOCH₃ and OH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 41.7, 63.6, 114.2, 117.6, 119.3, 128.2, 140.0, 159.4, 166.1.

4-Glutarylaminomethylsalicyl(O-methyl)hydroxamic Acid Succinimidyl Ester.

4-Aminomethylsalicyl(*O*-methyl)hydroxamic acid hydrochloride (0.57 g, 1.82 mmoles) was dissolved in anhydrous *N,N*-dimethylformamide (10 mL), and *N,N*-diisopropylethyl amine (0.35 mL, 2.01 mmoles) and glutaric anhydride (0.23 g, 2.02 mmoles) were added. The mixture was stirred at room temperature for 26 hours, and then *N*-hydroxysuccinimide (0.23 g, 2.00 mmoles) and 1,3-dicyclohexylcarbodiimide (0.42 g, 2.01 mmoles) were added. The reaction mixture was stirred for an additional 24 hours at


-69-

room temperature. The mixture was then filtered and diluted with ethyl acetate (100 mL). The ethyl acetate solution was washed with 1 N aqueous hydrochloric acid (100 mL), water (100 mL), and saturated aqueous sodium chloride (50 mL). The solution was then dried over anhydrous magnesium sulfate, filtered, and evaporated to dryness. The crude gummy product was triturated under a mixture of ethyl acetate and diethyl ether to produce a solid which was filtered and dried *in vacuo* to afford 0.15 g (20% yield) of 4-glutarylaminomethylsalicyl(*O*-methyl)hydroxamic acid succinimidyl ester (m.p. 121-124° C, open capillary, uncorrected).

¹H NMR (300 MHz, DMSO-*d*₆) δ 1.85 (quintet, *J* = 7 Hz, 2H, CH₂CH₂CH₂), 2.27 (triplet, *J* = 7 Hz, 2H, CH₂CONHCH₂), 2.69 (triplet, *J* = 7 Hz, 2H, NOCCH₂), 2.80 (singlet, 4H, COCH₂CH₂CO), 3.70 (singlet, 3H, NHOCH₃), 4.21 (doublet, *J* = 6 Hz, 2H, NHCH₂), 6.75 (doublet, *J* = 8 Hz, 1H, ArH), 6.77 (singlet, 1H, ArH), 7.57 (doublet, *J* = 8 Hz, 1H, ArH), 8.39 (triplet, *J* = 6 Hz, 1H, NHCH₂), 11.75 (singlet, 1H, OH), 11.78 (singlet, 1H, NHOCH₃). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 20.4, 25.4, 29.7, 33.4, 33.6, 41.7, 112.8, 115.6, 117.8, 127.8, 146.2, 159.3, 166.3, 169.0, 170.5, 171.5.

Example XI

Preparation Of A Reagent Of General Formula III 4-(3-(2-Pyridyldithio)propionyl)aminomethylsalicyl(*O*-methyl)hydroxamic Acid



-70-

4-(3-(2-Pyridyldithio)propionyl)aminomethylsalicyl(O-methyl)hydroxamic Acid.

3-(2-Pyridyldithio)propionic acid *N*-hydroxysuccinimide ester (100 mg, 0.32 mmole) was dissolved in dry *N,N*-dimethylformamide (5.0 mL) and *N,N*-diisopropylethylamine (65 μ L, 0.36 mmole) was added followed by 4-aminomethylsalicyl(*O*-methyl)hydroxamic acid hydrochloride (82 mg, 0.35 mmole). The reaction was stirred for 6 hours at room temperature. The solvent was evaporated *in vacuo*, and the residue was chromatographed on silica gel eluting with dichloromethane/methanol/acetic acid (95:5:1 [v/v/v]). Fractions containing the desired product were pooled and evaporated to an oil to afford 44 mg (35% yield) of 4-(3-(2-pyridyldithio)-propionyl)aminomethylsalicyl(*O*-methyl)hydroxamic acid.


^1H NMR (300 MHz, DMSO- d_6) δ 2.58 (triplet, $J = 7$ Hz, 2H, $\text{CH}_2\text{CH}_2\text{S}$), 3.04 (triplet, $J = 7$ Hz, 2H, COCH_2CH_2), 3.70 (singlet, 3H, OCH_3), 4.23 (doublet, $J = 6$ Hz, 2H, ArCH_2NH), 6.76 (doublet, $J = 8$ Hz, 1H, ArH), 6.79 (singlet, 1H, ArH), 7.23 (triplet, $J = 6$ Hz, 1H, ArH), 7.57 (doublet, $J = 8$ Hz, 1H, ArH), 7.73-7.82 (multiplet, 2H, ArH), 8.44 (doublet, $J = 6$ Hz, 1H, ArH), 8.48 (triplet, $J = 6$ Hz, 1H, NH), 11.80 (broad singlet, 2H, OH and NHO). ^{13}C NMR (75 MHz, DMSO- d_6) δ 34.1, 34.5, 41.8, 63.5, 112.7, 115.7, 117.8, 119.3, 121.3, 127.7, 138.0, 145.9, 149.8, 159.4, 166.3, 170.2, 172.8.

**PHENYLBORONIC ACID COMPLEXING REAGENTS AND
CONJUGATES
DERIVED FROM 4-AMINOMETHYL-2,6-DIHYDROXYBENZOIC
ACID**

A three-step process which utilizes reagents of General Formula CI for the preparation of bioconjugates is summarized in *Figure 1'*. Initially, a reagent of

-71-

General Formula CI is selected that is comprised of an appropriate reactive electrophilic or nucleophilic group R_1 suitable for reaction with the desired biologically active species.



General Formula CI


Group R_1 is a reactive electrophilic or nucleophilic moiety suitable for reaction of the putative phenylboronic acid complexing reagent with a bioactive species. Group R_1 is preferably selected from, but not limited to, acrylamide, bromo, dithiopyridyl, bromoacetamide, hydrazide, *N*-hydroxysuccinimidyl ester, *N*-hydroxysulfosuccinimidyl ester, imidate ester, imadazolidine, iodo, iodoacetamide, maleimide, amino and thiol moieties.

Group Z is a spacer selected from a saturated or unsaturated, preferably unbranched, chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide or disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length. Group Z is preferably selected from an unbranched alkyl chain of general formula $(CH_2)_n$, wherein $n = 1$ to 6.

Group R_2 is selected from an alkyl (e.g., methyl, ethyl, etc.) and a methylene bearing an electronegative substituent. An electronegative substituent is a substituent with a negative dipole moment, e.g., CN, COOH, etc. Group R_2 is preferably selected from one of CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 and CH_2OCH_3 .

The next step in a three-step process in the preparation of bioconjugates is to condense the appropriate reagent with the bioactive species to yield a conjugate of General Formula CII:


-72-



General Formula CII

In General Formula CII, Z and R₂ are as defined above, and BAS represents a biologically active species which may or may not contain a portion of a reactive moiety (and any spacer) used to attach the biologically active species to the reagent.


Next, the conjugate is reacted with a hydroxylamine derivative of the general formula NH₂OR₃, wherein R₃ is selected from either an H, an alkyl (e.g., methyl, ethyl, etc.), or a methylene or ethylene with an electronegative moiety. Suitable hydroxylamine derivatives include, but not limited to, NH₂OH, NH₂OCH₃, NH₂OCH₂CN, NH₂OCH₂COOH, NH₂OCH₂CONH₂ and NH₂OCH₂CH₂OH. The resulting conjugate has the general formula of General Formula CIV:



General Formula CIV

In General Formula CIV, Groups Z and BAS are as defined above, and R₃ is selected from one of an H, an alkyl, and a methylene moiety with an electronegative substituent.


The conjugate of General Formula CIV is then complexed with a phenylboronic acid having the general formula of General Formula CV:



General Formula CV

-73-


wherein the symbol labeled *BAS** represents a second biologically active species, that may include a linker portion and differ from the biologically active species labeled *BAS* of the complexing reagent. The *BAS** may also include a portion of a reactive moiety used to attach the bioactive species to the phenylboronic acid reagent. The complexation yields the stereoisomeric complex (about tetrahedral boron) of General Formula CVI:



General Formula CVI

It will be appreciated by one of ordinary skill in the art, upon reference to this disclosure, that the various syntheses described above in the previous section (concerning reagents and conjugates derived from 4 or 5 aminomethyl-salicylic acid and shown in Figures 1 through 7) may be utilized where appropriate for the corresponding synthesis of the reagents shown in Figures 1'-7' and/or described in this section. For example, the techniques for adding Z groups and R1 groups from the previous section may be considered for use in preparing the reagents shown in Figures 1' through 7' and/or described in this section.

SYNTHESIS OF PUTATIVE PHENYLBORONIC ACID COMPLEXING REAGENTS OF GENERAL FORMULA CI



General Formula CI

Reagents of General Formula CI are derived from alkyl 2,6-dihydroxy-4-methyl-benzoates. **Figure 2'** summarizes the preparation of alkyl 4-aminomethyl-2,6-dihydroxy-benzoates, synthetic intermediates leading to reagents of General Formula CI, wherein R_2 is an alkyl group, e.g., methyl, ethyl, etc. **Figure 2'** shows an example where R_2 is a methyl group. Initially, in step 1, an alkyl 2,6-dihydroxy-4-methylbenzoate is acetylated to reduce the reactivity of the ring system toward bromination, affording the corresponding alkyl 2,6-diacetoxy-4-methylbenzoate. In step 2, the ester is brominated with *N*-bromosuccinimide and benzoyl peroxide catalyst to afford the corresponding benzyl bromide. In step 3, the benzyl bromide is alkylated with sodium azide to afford the corresponding benzyl azide. In step 4, the benzyl azide is subjected to palladium catalyzed hydrogenation in the presence of HCl to afford the corresponding benzyl amine hydrochloride. Finally, in step 5, the acetoxy protecting groups are removed by acid catalyzed methanolysis.

Figure 3' summarizes the preparation of synthetic intermediates leading to reagents of General Formula CI, wherein group R_2 is a methylene bearing an electronegative moiety, for example, carboxymethyl, acetamidomethyl and cyanomethyl 4-aminomethyl-2,6-dihydroxy-benzoate. It is to be appreciated that other contemplated substituents for group R_2 may be prepared by a similar synthesis. Initially, in step 1, the alkyl 4-aminomethyl-2,6-dihydroxy-benzoate, prepared as summarized in **Figure 2'**, is reacted with di-*tert*-butyl dicarbonate and triethylamine in methanol to afford the corresponding protected alkyl *N*-(*tert*-butoxy-carbonyl)aminomethylbenzoate. In step 2, the alkyl ester is cleaved by reaction with potassium trimethylsilanolate and worked up in aqueous acid to afford the corresponding benzoic acid. In step 3, the benzoic acid is alkylated by reaction with either an α -haloacid, α -haloacetamide or α -haloacetonitrile and triethylamine to afford the corresponding carboxymethyl, acetamido-methyl or cyanomethyl ester, respectively. Finally, in step 4, the *N*-(*tert*-butoxycarbonyl) protecting group is removed by reaction with anhydrous HCl in tetrahydrofuran to afford an alkyl 4-aminomethyl-2, 6-dihydroxybenzoate hydrochloride.

Another reagent of the present invention, which is the acrylic acid amide of alkyl 4-amino methyl 2, 6 dihydroxy-benzoate can be prepared in a single step, using the end product of **Figure 3** by condensing acrylic acid anhydride or acryloyl chloride with alkyl 4- aminomethyl 2, 6 dihydroxy-benzoate.

-75-

Figure 5' summarizes the synthesis of reagents of General Formula CI, wherein group R_2 is one of either an alkyl or a methylene bearing an electronegative moiety, and wherein group R_1 is selected from either imidazolidine, hydrazide and *N*-hydroxysuccinimidyl ester moieties. These reagents are each prepared by a two-step process in which an aliphatic acid anhydride is utilized in the first step. Initially, an alkyl 4-aminomethyl-2,6-dihydroxybenzoate, prepared as summarized in either *Figure 2'* or *Figure 3'*, is condensed with an aliphatic acid anhydride preferably selected from, but not limited to, either succinic anhydride, glutaric anhydride, maleic anhydride and glycolic acid anhydride, in an aprotic organic solvent, which results in the introduction of a spacer (group Z) having a free terminal carboxylic acid moiety. In the case where the aliphatic acid anhydride is maleic anhydride in this condensation reaction, the resulting Z group is unsaturated as it contains an alkene group. Subsequently, the carboxylic acid moiety is further functionalized by reaction with either *N,N*-carbonyldiimidazole, isobutylchloroformate and *tert*-butyl carbazate, or *N,N*-dicyclohexylcarbodiimide and *N*-hydroxy-succinimide to afford the corresponding imidazolidine, protected hydrazide and *N*-hydroxy-succinimidyl ester, respectively. In the instance of the protected hydrazide, the *N*-(*tert*-butoxycarbonyl) protecting group is removed by contacting the reagent with anhydrous hydrochloric acid.

Figure 6' summarizes the synthesis of reagents of General Formula CI, wherein group R_2 is one of either an alkyl or a methylene bearing an electronegative moiety, and wherein group R_1 is selected from either acrylamide, bromoacetamide, chloroacetamide, maleimide, dithiopyridyl and imidate ester moieties. Reagents of General Formula CI, wherein group R_1 is selected from either bromo and chloro moieties, are prepared by condensing an alkyl 4- or 5-aminomethyl-salicylate, prepared as summarized in either *Figure 2'* or *Figure 3'*, with either bromoacetic acid anhydride or chloroacetic acid anhydride, respectively. The homologous iodo reagent is prepared by halogen exchange of the chloro reagent with sodium iodide. Reagents of General Formula CI, wherein R_1 is selected from either bromo, chloro, iodo, bromoacetamide, chloroacetamide and iodoacetamide moieties, may not be conveniently prepared when R_3 is H, due to the potential for intermolecular alkylation of the unprotected hydroxamate. Reagents of General Formula CI, wherein group R_1 is selected from either maleimide and dithiopyridyl moieties, are prepared by condensing an alkyl 4-

aminomethyl-2,6-dihydroxybenzoate, prepared as summarized in either *Figure 2'* or *Figure 3'*, with an *N*-hydroxysuccinimidyl ester of an aliphatic carboxylic ester which bears either a terminal maleimide or dithiopyridyl moiety. Reagents of General Formula CI, wherein R_1 is an imidate ester moiety, are prepared by a two-step process in which an alkyl 4-aminomethyl-2,6-dihydroxybenzoate, prepared as summarized in either *Figure 2'* or *Figure 3'*, is first condensed with an *N*-hydroxysuccinimidyl ester of an aliphatic carboxylic ester which bears a terminal nitrile moiety. Subsequently, the nitrile moiety is converted to the methyl imidate ester by reaction with anhydrous hydrogen chloride acid in methanol at 0° C.

Reagents of General Formula CI, wherein group R_1 is selected from either *N*-hydroxy-succinimidyl ester and dithiopyridyl moieties, may be utilized as synthetic intermediates to prepare reagents of General Formula CI, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide or disulfide moieties.

Reagents of General Formula CI, wherein group R_1 is an *N*-hydroxysuccinimidyl ester moiety, may be condensed with compounds having primary aliphatic amine moieties of the general formula $R_1-Z_2-NH_2$, wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford reagents of General Formula CI, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide moieties.

Alternatively, *N*-hydroxysuccinimidyl ester reagents of General Formula CI, preferably derived from a dicarboxylic acid selected from either succinic acid, maleic acid, fumaric acid, acetylenedicarboxylic acid and glutaric acid, may be condensed with compounds having primary aliphatic amine moieties of the general formula $HO_2C-Z_2-NH_2$, wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, preferably selected from, but not limited to, either glycine, β -alanine, aminopropionic acid, 4-aminobutyric acid and 6-aminocaproic acid, to afford compounds having free terminal carboxylic acid moieties which may be further functionalized in accordance with *Figure 5'* to afford reagents of General Formula CI, wherein Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide moieties. This process is summarized in *Figure 7'* for the synthesis of a reagent of General Formula CI, wherein R_1 is an *N*-hydroxysuccinimidyl ester,

-77-

and wherein Z is an unbranched saturated or unsaturated chain with at least one of an intermediate amide moiety.


Reagents of General Formula CI, wherein group R_1 is a dithiopyridyl moiety, may be condensed with compounds having terminal thiol moieties of the general formula R_1-Z_2-SH , wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford reagents of General Formula CI, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate disulfide moieties.

Alternatively, dithiopyrinyll reagents of General Formula CI, preferably derived from a mercaptocarboxylic acid selected from either mercaptoacetic acid, β -mercaptopropionic acid, mercaptopropiolic acid, 4-mercaptobutyric acid and 6-mercaptocaproic acid, may be condensed with compounds having a thiol moiety of the general formula HO_2C-Z_2-SH , wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, preferably selected from, either mercaptoacetic acid, β -mercaptopropionic acid, mercaptopropiolic acid, 4-mercaptobutyric acid and 6-mercaptocaproic acid, to afford compounds having free terminal carboxylic acid moieties which may be further functionalized in accordance with *Figure 5'* to afford reagents of General Formula I, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate disulfide moieties. This process is summarized in *Figure 7* for the synthesis of a reagent of General Formula CI, wherein R_1 is an *N*-hydroxysuccinimidyl ester, and wherein Z is an unbranched saturated or unsaturated chain with at least one of an intermediate disulfide moiety.

Reagents of General Formula CI, wherein group Z is a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length, are prepared by condensing an alkyl 4-aminomethyl-2,6-dihydroxybenzoate, prepared as summarized in either *Figure 2'* or *Figure 3'*, with a polyethylene glycol reagent having both an *N*-hydroxysuccinimidyl ester moiety and either a reactive electrophilic or nucleophilic moiety (or a protected precursor thereof), many of which are commercially available, to afford reagents of General Formula CI, wherein group Z is a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length.

-78-

SYNTHESIS OF
PHENYLBORONIC ACID COMPLEXING REAGENTS
OF GENERAL FORMULA CIII




General Formula CIII

Figure 4' summarizes the preparation of 4-aminomethyl-2,6-dihydroxybenzohydroxamic acids, synthetic intermediates leading to reagents of General Formula CIII, wherein group R_3 is one of an alkyl or methylene bearing an electronegative moiety. Initially, in step 1, an alkyl 4-aminomethyl-2,6-dihydroxybenzoate, prepared as summarized in *Figure 2'*, is condensed with *N*-(benzyloxycarbonyl)oxy succinimide to afford the alkyl *N*-benzyloxycarbonyl protected alkyl 4-aminomethyl-2,6-dihydroxybenzoate. In step 2, the phenolic hydroxyl moieties are condensed with benzyl bromide to afford the further protected benzyl ether intermediate. In step 3, the alkyl ester is selectively cleaved by reaction with LiOH to afford the corresponding carboxylic acid. In step 4, the carboxylic acid is activated by reaction with isobutylchloroformate to form a mixed anhydride which is subsequently reacted with a hydroxylamine derivative preferably selected from, but not limited to, either NH_2OH , NH_2OCH_3 , $\text{NH}_2\text{OCH}_2\text{CN}$, $\text{NH}_2\text{OCH}_2\text{COOH}$, $\text{NH}_2\text{OCH}_2\text{CONH}_2$, $\text{NH}_2\text{OCH}_2\text{CH}_2\text{OH}$ and $\text{NH}_2\text{OCH}_2\text{OCH}_3$ to afford the corresponding protected hydroxamic acid. Finally, in step 5, both the amine and phenolic hydroxyl moieties are simultaneously deprotected by palladium catalyzed hydrogenation in the presence of HCl to afford the corresponding 4-aminomethyl-2,6-dihydroxybenzohydroxamic acid hydrochloride. These synthetic intermediates, which are structurally related to the corresponding alkyl 4-amino-methyl-2,6-dihydroxybenzoates, may be further functionalized in accordance with either *Figure 5'* or *Figure 6'* to afford reagents of General Formula CIII.

-79-


PREPARATION OF
PHENYLBORONIC ACID COMPLEXING CONJUGATES
OF GENERAL FORMULA CIV

At this point, the putative phenylboronic acid complexing reagents of General Formula CI may be reacted with a suitable biologically active species to yield the conjugate of the general formula of General Formula CII:



General Formula CII

The conjugate of General Formula CII is next condensed with a hydroxylamine derivative to yield the corresponding phenylboronic acid complexing conjugate of the general formula of General Formula CIV:




General Formula CIV

Suitable hydroxylamine derivatives are preferably selected from, but not limited to, NH_2OH , NH_2OCH_3 , $\text{NH}_2\text{OCH}_2\text{CN}$, $\text{NH}_2\text{OCH}_2\text{COOH}$, $\text{NH}_2\text{OCH}_2\text{CONH}_2$, $\text{NH}_2\text{OCH}_2\text{CH}_2\text{OH}$, $\text{NH}_2\text{OCH}_2\text{OCH}_3$. When group R_2 in General Formula CII is an alkyl group, NH_2OH is preferably utilized to effect the interconversion of General Formula CII to General Formula CIV.


Alternatively, conjugates of General Formula CIV may also be prepared by the route which is also described in *Figure 1'* which utilizes a reagent of General Formula CIII:

-80-




General Formula CIII

Phenylboronic acid complexing reagents of General Formula CIII may be reacted with a suitable biologically active species to yield the conjugate of the general formula of General Formula CIV:



General Formula CIV

A two-step process which utilizes reagents of General Formula CIII for the preparation of bioconjugates is summarized in *Figure 1'*. Initially, a reagent of General Formula CIII is selected that is comprised of an appropriate reactive electrophilic or nucleophilic group R_1 suitable for reaction with the desired biologically active species.



General Formula CIII

Group R_1 is a reactive electrophilic or nucleophilic moiety suitable for reaction of the phenylboronic acid complexing reagent with a bioactive species. Group R_1 is preferably selected from, but not limited to, acrylamide, bromo, dithiopyridyl, bromoacetamide, hydrazide, *N*-hydroxysuccinimidyl ester, *N*-hydroxysulfosuccinimidyl ester, imidate ester, imidazolidine, iodo, iodoacetamide, maleimide, amino and thiol moieties.


-81-

Group Z is a spacer selected from a saturated or unsaturated, preferably unbranched, chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide or disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length. Group Z is preferably selected from an unbranched alkyl chain of general formula $(CH_2)_n$, wherein $n = 1$ to 6.

Group R_3 is selected from one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent.

Group R_3 is preferably selected from one of H, CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 , CH_2CH_2OH and CH_2OCH_3 . When group R_3 is H, group R_1 is preferably selected from one of acrylamide, amino, dithiopyridyl, hydrazide, imidate ester, maleimide, and thiol moieties.


The reagent of General Formula CIII is condensed with the bioactive species to yield a conjugate of General Formula CIV:



General Formula CIV

wherein groups Z and BAS are as defined above, and R_3 is selected from one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent.

The conjugate of General Formula CIV is then complexed with a phenylboronic acid conjugate having the general formula of General Formula CV:




General Formula CV

wherein the symbol labeled BAS^* represents a second biologically active species, that may include a linker portion and differ from the biologically active species


-82-

labeled *BAS* of the complexing reagent. The *BAS** may also include a portion of a reactive moiety used to attach the bioactive species to the phenylboronic acid reagent. The complexation yields the stereoisomeric complex (tetrahedral boron) of General Formula CVI:



General Formula CVI

An alternative three-step process which utilizes reagents of General Formula CI for the preparation of bioconjugates is also summarized in *Figure 1'*. Initially, a reagent of General Formula CI is selected that is comprised of an appropriate reactive electrophilic or nucleophilic group R_1 suitable for reaction with the desired biologically active species.



General Formula CI

Group R_1 is a reactive electrophilic or nucleophilic moiety suitable for reaction of the putative phenylboronic acid complexing reagent with a bioactive species. Group R_1 is preferably selected from, but not limited to, acrylamide, bromo, dithiopyridyl, bromoacetamide, hydrazide, *N*-hydroxysuccinimidyl ester, *N*-hydroxysulfosuccinimidyl ester, imidate ester, imidazolidine, iodo, iodoacetamide, maleimide, amino and thiol moieties.


Group Z is a spacer selected from a saturated or unsaturated, preferably unbranched, chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide or disulfide moieties,

-83-

and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length. Group Z is preferably selected from an unbranched alkyl chain of general formula $(CH_2)_n$, wherein $n = 1$ to 6.

Group R_2 is selected from an alkyl (e.g., methyl, ethyl, etc.) and a methylene or ethylene bearing an electronegative substituent. An electronegative substituent is a substituent with a negative dipole moment, e.g., CN, COOH, etc. Group R_2 is preferably selected from one of CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 and CH_2OCH_3 .


The next step in a three-step process in the preparation of bioconjugates is to condense the appropriate reagent with the bioactive species to yield a conjugate of General Formula CII:



General Formula CII

In General Formula CII, Z, BAS and R_2 are as defined above.

Next, the conjugate is reacted with a hydroxylamine derivative of the general formula NH_2OR_3 , wherein R_3 is selected from either an H, an alkyl (e.g., methyl, ethyl, etc.), or a methylene or ethylene with an electronegative moiety. Suitable hydroxylamine derivatives include, but not limited to, NH_2OH , NH_2OCH_3 , NH_2OCH_2CN , NH_2OCH_2COOH , $NH_2OCH_2CONH_2$ and $NH_2OCH_2CH_2OH$. The resulting conjugate has the general formula of General Formula CIV:




General Formula CIV

-84-


In General Formula CIV, Groups *Z* and *BAS* are as defined above, and *R*₃ is selected from one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent.

The conjugate of General Formula CIV is then complexed with a phenylboronic acid having the general formula of General Formula CV:



General Formula CV

wherein the symbol labeled *BAS** represents a second biologically active species, that may include a linker portion and differ from the biologically active species labeled *BAS* of the complexing reagent. The *BAS** may also include a portion of a reactive moiety used to attach the bioactive species to the phenylboronic acid reagent. The complexation yields the stereoisomeric complex (about tetrahedral boron) of General Formula CVI:




General Formula CVI

It will be appreciated by one of the ordinary skill in the art, upon reference to this disclosure, that the various syntheses described above in the previous section (concerning reagents and conjugates derived from 4 or 5 aminomethyl-salicylic acid and shown in Figures 1 through 7) may be utilized where appropriate for the corresponding synthesis of the reagents shown in Figures 1'-7' and/or described in this section. For example, the techniques for adding *Z* groups and *R*₁ group from the previous section may be considered for use in preparing the reagents shown in Figures 1'-7' and/or described in this section.

-85-

SYNTHESIS OF
PHENYLBORONIC ACID COMPLEXING REAGENTS
OF GENERAL FORMULA CIII



General Formula CIII

Reagents of General Formula CIII are derived from 2,6-dihydroxy-4-methylbenzoic acid. In each instance, the reagent is ultimately prepared from a synthetic intermediate which is an alkyl 4-aminomethyl-2,6-dihydroxybenzoate.

Figure 2' summarizes the preparation of alkyl 4-aminomethyl-2,6-dihydroxybenzoates, synthetic intermediates leading to reagents of General Formula CI, wherein R_2 is an alkyl group, e.g., methyl, ethyl, etc. Initially, in step 1, an alkyl 2,6-dihydroxy-4-methylbenzoate is acetylated to reduce the reactivity of the ring system toward bromination, affording the corresponding alkyl 2,6-diacetoxy-4-methylbenzoate. In step 2, the ester is brominated with *N*-bromosuccinimide and benzoyl peroxide catalyst to afford the corresponding benzyl bromide. In step 3, the benzyl bromide is alkylated with sodium azide to afford the corresponding benzyl azide. In step 4, the benzyl azide is subjected to palladium catalyzed hydrogenation in the presence of HCl to afford the corresponding benzyl amine hydrochloride. Finally, in step 5, the acetoxy protecting groups are removed by acid catalyzed methanolysis.

Figure 4' summarizes the preparation of 4-aminomethyl-2,6-dihydroxybenzohydroxamic acids, synthetic intermediates leading to reagents of General Formula CIII, wherein group R_3 is one of an alkyl or methylene or ethylene bearing an electronegative substituent. Initially, in step 1, an alkyl 4-aminomethyl-2,6-dihydroxybenzoate, prepared as summarized in *Figure 2'*, is condensed with *N*-(benzyloxycarbonyl)oxy succinimide to afford the alkyl *N*-benzyloxycarbonyl protected 4-aminomethyl-2,6-dihydroxybenzoate. In step 2, the phenolic hydroxyl moieties are condensed with benzyl bromide to afford the further protected benzyl diether intermediate. In step 3, the alkyl ester is

selectively cleaved by reaction with LiOH to afford the corresponding benzoic acid. In step 4, the benzoic acid is activated by reaction with isobutylchloroformate to form a mixed anhydride which is subsequently reacted with a hydroxylamine derivative preferably selected from, but not limited to, either NH_2OH , NH_2OCH_3 , $\text{NH}_2\text{OCH}_2\text{CN}$, $\text{NH}_2\text{OCH}_2\text{COOH}$, $\text{NH}_2\text{OCH}_2\text{CONH}_2$, $\text{NH}_2\text{OCH}_2\text{CH}_2\text{OH}$ and $\text{NH}_2\text{OCH}_2\text{OCH}_3$ to afford the corresponding protected hydroxamic acid. Finally, in step 5, both the amine and phenolic hydroxyl moieties are simultaneously deprotected by palladium catalyzed hydrogenation in the presence of HCl to afford the corresponding 4-aminomethyl-2,6-dihydroxybenzohydroxamic acid hydrochloride.

Another reagent of the present invention, which is the acrylic acid amide of 4-aminomethyl 2, 6 dihydroxybenzo hydroxamic acid, can be prepared in a single step, using the end product of Figure 4', by condensing acrylic acid anhydride or acryloyl chloride with 4 aminomethyl 2, 6 dihydroxybenzo hydroxamic acid, provided that R_3 is not H.

Figure 5' summarizes the synthesis of reagents of General Formula CIII, wherein group R_3 is preferably selected from one of H, CH_3 , and a methylene or ethylene moiety with an electronegative substituent, and wherein group R_1 is selected from either imidazolidine, hydrazide and *N*-hydroxysuccinimidyl ester moieties. These reagents are each prepared by a two-step process in which an aliphatic acid anhydride is utilized in the first step. Initially, a 4-amino-methyl-2,6-dihydroxybenzohydroxamic acid, prepared as summarized in **Figure 4'**, is condensed with an aliphatic acid anhydride preferably selected from, but not limited to, either succinic anhydride, glutaric anhydride, maleic anhydride and glycolic acid anhydride, in an aprotic organic solvent, which results in the introduction of a spacer (group Z) having a free terminal carboxylic acid moiety. In the case where the aliphatic acid anhydride is maleic anhydride in this condensation reaction, the resulting Z group is unsaturated as it contains an alkene group. Subsequently, the carboxylic acid moiety is further functionalized by reaction with either *N,N*-carbonyldiimidazole, isobutylchloroformate and *tert*-butyl carbazate, or *N,N*-dicyclohexyl-carbodiimide and *N*-hydroxysuccinimide to afford the corresponding imidazolidine, protected hydrazide and *N*-hydroxysuccinimidyl ester, respectively. In the instance of the protected hydrazide, the *N*-(*tert*-butoxycarbonyl) protecting group is removed by contacting the reagent with anhydrous hydrochloric acid.

Figure 6' summarizes the synthesis of reagents of General Formula CIII, wherein group R_3 is preferably selected from one of H, CH_3 , and a methylene or ethylene moiety with an electronegative substituent, and wherein group R_1 is selected from either bromo, chloro, maleimide, dithiopyridyl and imidate ester moieties. Reagents of General Formula CIII, wherein group R_1 is selected from either bromo and chloro moieties, are prepared by condensing a 4-aminomethyl-2,6-dihydroxybenzohydroxamic acid, prepared as summarized in *Figure 4'*, with either bromoacetic acid anhydride or chloroacetic acid anhydride, respectively. The homologous iodo reagent is prepared by halogen exchange of the chloro reagent with sodium iodide. Reagents of General Formula CIII, wherein R_1 is selected from either bromo, chloro, iodo, bromoacetamide, chloroacetamide and iodoacetamide moieties, may not be conveniently prepared when R_3 is H, due to the potential for intermolecular alkylation of the unprotected hydroxamate. Reagents of General Formula CIII, wherein group R_1 is selected from either maleimide and dithiopyridyl moieties, are prepared by condensing a 4-aminomethyl-2,6-dihydroxybenzohydroxamic acid, prepared as summarized in *Figure 4'*, with an *N*-hydroxysuccinimidyl ester of an aliphatic carboxylic ester which bears either a terminal maleimide or dithiopyridyl moiety. Reagents of General Formula CIII, wherein R_1 is an imidate ester moiety, are prepared by a two-step process in which a 4-aminomethyl-2,6-dihydroxybenzohydroxamic acid, prepared as summarized in *Figure 4'*, is first condensed with an *N*-hydroxysuccinimidyl ester of an aliphatic carboxylic ester which bears a terminal nitrile moiety. Subsequently, the nitrile moiety is converted to the methyl imidate ester by reaction with anhydrous hydrogen chloride in methanol at $0^\circ C$.

Reagents of General Formula CIII, wherein group R_1 is selected from either *N*-hydroxy-succinimidyl ester and dithiopyridyl moieties may be utilized as synthetic intermediates to prepare reagents of General Formula CIII, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide or disulfide moieties.

Reagents of General Formula CIII, wherein group R_1 is an *N*-hydroxysuccinimidyl ester moieties, may be condensed with compounds having primary aliphatic amine moieties of the general formula $R_1-Z_2-NH_2$, wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford reagents of General Formula CIII, wherein

-88-

group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide moieties.

Alternatively, *N*-hydroxysuccinimidyl ester reagents of General Formula CIII, preferably derived from a dicarboxylic acid selected from either succinic acid, maleic acid, fumaric acid, acetylenedicarboxylic acid and glutaric acid, may be condensed with compounds having primary aliphatic amine moieties of the general formula $\text{HO}_2\text{C}-\text{Z}_2-\text{NH}_2$, wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, preferably selected from, either glycine, β -alanine, α -aminopropionic acid, 4-amino-butyric acid and 6-aminocaproic acid, to afford compounds having free terminal carboxylic acid moieties which may be further functionalized in accordance with *Figure 5'* to afford reagents of General Formula CIII, wherein Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide moieties. This process is summarized for in *Figure 7'* for the synthesis of a reagent of General Formula CIII, wherein R_1 is an *N*-hydroxysuccinimidyl ester, and wherein Z is an unbranched saturated or unsaturated chain with at least one of an intermediate amide moiety.

Reagents of General Formula CIII, wherein group R_1 is a dithiopyridyl moiety, may be condensed with compounds having terminal thiol moieties of the general formula $\text{R}_1-\text{Z}_2-\text{SH}$, wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford reagents of General Formula CIII, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate disulfide moieties.


Alternatively, dithiopyridyl reagents of General Formula CIII, preferably derived from a mercaptocarboxylic acid selected from either mercaptoacetic acid, β -mercaptopropionic acid, mercaptopropionic acid, 4-mercaptobutyric acid and 6-mercaptocaproic acid, may be condensed with compounds having a terminal thiol moiety of the general formula $\text{HO}_2\text{C}-\text{Z}_2-\text{SH}$, wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, preferably selected from either mercaptoacetic acid, β -mercaptopropionic acid, mercaptopropionic acid, 4-mercaptobutyric acid and 6-mercaptocaproic acid, to afford compounds having free terminal carboxylic acid moieties which may be further functionalized in accordance with *Figure 5'* to afford reagents of General Formula CIII, wherein group Z is an unbranched saturated or unsaturated chain

-89-

with at least one of intermediate disulfide moieties. This process is summarized in *Figure 7'* for the synthesis of a reagent of General Formula CIII, wherein R_1 is an *N*-hydroxysuccinimidyl ester, and wherein Z is an unbranched saturated or unsaturated chain with at least one of a intermediate disulfide moiety.

Reagents of General Formula CIII, wherein group Z is a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length, are prepared by condensing a 4-aminomethyl-2,6-dihydroxybenzohydroxamic acid, prepared as summarized in *Figure 4'*, with a polyethylene glycol reagent having both an *N*-hydroxysuccinimidyl ester moiety and either a reactive electrophilic or nucleophilic moiety (or a protected precursor thereof), many of which are commercially available, to afford reagents of General Formula CIII, wherein group Z is a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length.

SYNTHESIS OF PUTATIVE PHENYLBORONIC ACID COMPLEXING REAGENTS OF GENERAL FORMULA CI



General Formula CI

Reagents of General Formula CI are derived from alkyl 2,6-dihydroxy-4-methylbenzoates. *Figure 2'* summarizes the preparation of alkyl 4-aminomethyl-2,6-dihydroxybenzoates, synthetic intermediates leading to reagents of General Formula CI, wherein R_2 is an alkyl group, e.g., methyl, ethyl, etc. *Figure 2'* shows an example where R_2 is a methyl group. Initially, in step 1, an alkyl 2,6-dihydroxy-4-methylbenzoate is acetylated to reduce the reactivity of the ring system toward bromination, affording the corresponding alkyl 2,6-diacetoxy-4-methylbenzoate. In step 2, the ester is brominated with *N*-bromosuccinimide and benzoyl peroxide catalyst to afford the corresponding benzyl bromide. In step

-90-

3, the benzyl bromide is alkylated with sodium azide to afford the corresponding benzyl azide. In step 4, the benzyl azide is subjected to palladium catalyzed hydrogenation in the presence of HCl to afford the corresponding benzyl amine hydrochloride. Finally, in step 5, the acetoxymethyl protecting groups are removed by acid catalyzed methanolysis.

Figure 3' summarizes the preparation of synthetic intermediates leading to reagents of General Formula CI, wherein group R_2 is a methylene bearing an electronegative substituent, for example, carboxymethyl, acetamidomethyl and cyanomethyl 4-aminomethyl-2,6-dihydroxybenzoate. It is to be appreciated that other contemplated substituents for group R_2 may be prepared by a similar synthesis. Initially, in step 1, the alkyl 4-aminomethyl-2,6-dihydroxybenzoate, prepared as summarized in **Figure 2'**, is reacted with di-*tert*-butyl dicarbonate and triethylamine in methanol to afford the corresponding protected alkyl *N*-(*tert*-butoxycarbonyl) aminomethylbenzoate. In step 2, the alkyl ester is cleaved by reaction with potassium trimethylsilanolate and worked up in aqueous acid to afford the corresponding benzoic acid. In step 3, the benzoic acid is alkylated by reaction with either an α -haloacid, α -haloacetamide or α -haloacetonitrile and triethylamine to afford the corresponding carboxymethyl, acetamido-methyl or cyanomethyl ester, respectively. Finally, in step 4, the *N*-(*tert*-butoxycarbonyl) protecting group is removed by reaction with anhydrous HCl in tetrahydrofuran to afford an alkyl 4-aminomethyl-2,6-dihydroxybenzoate hydrochloride.

Figure 5' summarizes the synthesis of reagents of General Formula CI, wherein group R_2 is one of either an alkyl or a methylene or ethylene bearing an electronegative substituent, and wherein group R_1 is selected from either imidazolide, hydrazide and *N*-hydroxysuccinimide ester moieties. These reagents are each prepared by a two-step process in which an aliphatic acid anhydride is utilized in the first step. Initially, an alkyl 4-aminomethyl-2,6-dihydroxybenzoate, prepared as summarized in either **Figure 2'** or **Figure 3'**, is condensed with an aliphatic acid anhydride preferably selected from, but not limited to, either succinic anhydride, glutaric anhydride, maleic anhydride and glycolic acid anhydride, in an aprotic organic solvent, which results in the introduction of a spacer (group Z) having a free terminal carboxylic acid moiety. Subsequently, the carboxylic acid moiety is further functionalized by reaction with either *N,N*-carbonyldiimidazole, isobutylchloroformate and *tert*-butyl carbazate, or *N,N*-dicyclohexylcarbodiimide

and *N*-hydroxysuccinimide to afford the corresponding imidazolidine, protected hydrazide and *N*-hydroxysuccinimidyl ester, respectively. In the instance of the protected hydrazide, the *N*-(*tert*-butoxycarbonyl) protecting group is removed by contacting the reagent with anhydrous hydrochloric acid.

Figure 6' summarizes the synthesis of reagents of General Formula CI, wherein group R_2 is one of either an alkyl or a methylene or ethylene bearing an electronegative substituent, and wherein group R_1 is selected from either bromo, chloro, maleimide, dithiopyridyl and imidate ester moieties. Reagents of General Formula CI, wherein group R_1 is selected from either bromo and chloro moieties, are prepared by condensing an alkyl 4-aminomethyl-2,6-dihydroxybenzoate, prepared as summarized in either **Figure 2'** or **Figure 3'**, with either bromoacetic acid anhydride or chloroacetic acid anhydride, respectively. The homologous iodo reagent is prepared by halogen exchange of the chloro reagent with sodium iodide. Reagents of General Formula CI, wherein R_1 is selected from either bromo, chloro, iodo, bromoacetamide, chloroacetamide and iodoacetamide moieties, may not be conveniently prepared when R_3 is H, due to the potential for intermolecular alkylation of the unprotected hydroxamate. Reagents of General Formula CI, wherein group R_1 is selected from either maleimide and dithiopyridyl moieties, are prepared by condensing an alkyl 4-aminomethyl-2,6-dihydroxybenzoate, prepared as summarized in either **Figure 2'** or **Figure 3'**, with an *N*-hydroxysuccinimidyl ester of an aliphatic carboxylic ester which bears either a terminal maleimide or dithiopyridyl moiety. Reagents of General Formula CI, wherein R_1 is an imidate ester moiety, are prepared by a two-step process in which an alkyl 4-aminomethyl-2,6-dihydroxybenzoate, prepared as summarized in either **Figure 2'** or **Figure 3'**, is first condensed with an *N*-hydroxysuccinimidyl ester of an aliphatic carboxylic ester which bears a terminal nitrile moiety. Subsequently, the nitrile moiety is converted to the methyl imidate ester by reaction with anhydrous hydrochloric acid in methanol at 0° C.

Reagents of General Formula CI, wherein group R_1 is selected from either dithiopyridyl, imidazolidine and *N*-hydroxysuccinimidyl ester moieties may be utilized as synthetic intermediates to prepare reagents of General Formula CI, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide or disulfide moieties.

Reagents of General Formula CI, wherein group R_1 is selected from either *N*-hydroxy-succinimidyl ester and dithiopyridyl moieties may be utilized as synthetic intermediates to prepare reagents of General Formula CI, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide or disulfide moieties.

Reagents of General Formula CI, wherein group R_1 is an *N*-hydroxysuccinimidyl ester moieties, may be condensed with compounds having primary aliphatic amine moieties of the general formula $R_1-Z_2-NH_2$, wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford reagents of General Formula CI, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide moieties.

Alternatively, *N*-hydroxysuccinimidyl ester reagents of General Formula CI, preferably derived from a dicarboxylic acid selected from either succinic acid, maleic acid, fumaric acid, acetylenedicarboxylic acid and glutaric acid, may be condensed with compounds having primary aliphatic amine moieties of the general formula $HO_2C-Z_2-NH_2$, wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, preferably selected from, either glycine, β -alanine, α -aminopropionic acid, 4-amino-butyric acid and 6-aminocaproic acid, to afford compounds having free terminal carboxylic acid moieties which may be further functionalized in accordance with *Figure 5'* to afford reagents of General Formula CI, wherein Z is an unbranched saturated or unsaturated chain with at least one of intermediate amide moieties. This process is summarized for in *Figure 7'* for the synthesis of a reagent of General Formula CI, wherein R_1 is an *N*-hydroxysuccinimidyl ester, and wherein Z is an unbranched saturated or unsaturated chain with at least one of an intermediate amide moiety.

Reagents of General Formula CI, wherein group R_1 is a dithiopyridyl moiety, may be condensed with compounds having terminal thiol moieties of the general formula R_1-Z_2-SH , wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, to afford reagents of General Formula CI, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate disulfide moieties.

Alternatively, dithiopyridyl reagents of General Formula CI, preferably

-93-


derived from a mercaptocarboxylic acid selected from either mercaptoacetic acid, β -mercaptopropionic acid, mercaptopropiolic acid, 4-mercaptobutyric acid and 6-mercaptocaproic acid, may be condensed with compounds having a terminal thiol moiety of the general formula $\text{HO}_2\text{C}-\text{Z}_2-\text{SH}$, wherein Z_2 is an unbranched saturated or unsaturated chain preferably of from about 1 to 5 carbon equivalents in length, preferably selected from either mercaptoacetic acid, β -mercaptopropionic acid, mercaptopropiolic acid, 4-mercaptobutyric acid and 6-mercaptocaproic acid, to afford compounds having free terminal carboxylic acid moieties which may be further functionalized in accordance with *Figure 5'* to afford reagents of General Formula CI, wherein group Z is an unbranched saturated or unsaturated chain with at least one of intermediate disulfide moieties. This process is summarized in *Figure 7'* for the synthesis of a reagent of General Formula CI, wherein R_1 is an *N*-hydroxysuccinimidyl ester, and wherein Z is an unbranched saturated or unsaturated chain with at least one of a intermediate disulfide moiety.

Reagents of General Formula CI, wherein group Z is a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length, are prepared by condensing an alkyl 4-aminomethyl-2,6-dihydroxybenzoate, prepared as summarized in either *Figure 2'* or *Figure 3'*, with a polyethylene glycol reagent having both an *N*-hydroxysuccinimidyl ester moiety and either a reactive electrophilic or nucleophilic moiety (or a protected precursor thereof), many of which are commercially available, to afford reagents of General Formula CI, wherein group Z is a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length.

PREPARATION OF PHENYLBORONIC ACID COMPLEXING CONJUGATES OF GENERAL FORMULA CIV


At this point, phenylboronic acid complexing reagents of General Formula CIII may be reacted with a suitable biologically active species to yield the conjugate of the general formula of General Formula CIV:

-94-




General Formula CIV

Alternatively, the putative phenylboronic acid complexing reagents of General Formula CI may be reacted with a suitable biologically active species to yield the conjugate of the general formula of General Formula CII:



General Formula CII

The conjugate of General Formula CII is next condensed with a hydroxylamine derivative to yield the corresponding phenylboronic acid complexing conjugate of the general formula of General Formula CIV:



General Formula CIV

Suitable hydroxylamine derivatives are preferably selected from, but not limited to, NH_2OH , NH_2OCH_3 , $\text{NH}_2\text{OCH}_2\text{CN}$, $\text{NH}_2\text{OCH}_2\text{COOH}$, $\text{NH}_2\text{OCH}_2\text{CONH}_2$, $\text{NH}_2\text{OCH}_2\text{CH}_2\text{OH}$, $\text{NH}_2\text{OCH}_2\text{OCH}_3$. When group R_2 in General Formula CII is an alkyl group, NH_2OH is preferably utilized to effect the interconversion of General Formula CII to General Formula CIV.

-95-

PREPARATION OF BIOCONJUGATES OF GENERAL FORMULA CVI

Bioconjugates of General Formula CVI may be prepared in buffered aqueous solutions or organic solvents. Preferred buffers include acetate, citrate, phosphate, carbonate and diglycine. Borate buffers should be avoided due to their ability to complex with the phenylboronic acid complexing moiety. Tris, β -hydroxyamine and β -hydroxyacid buffers should be avoided due to their ability to complex with the phenylboronic acid. The bioconjugate is formed within a few minutes over a range of temperatures of from about 4°C to 70°C. The stability of the bioconjugate in aqueous solution at a given pH and temperature is influenced, to some extent, by substituent group R_3 . Bioconjugates of General Formula CVI are stable in aqueous solutions of approximate pH greater than 2.5 and less than 12.5. The bioconjugation reaction (phenylboronic acid complexation) is insensitive to significant variations in ionic strength over the range 0.01 to 1 M, the presence of organic solvents including acetonitrile, methanol, ethanol, isopropanol, butanol, *N,N*-dimethylformamide and dimethylsulfoxide, the presence of detergents, and the presence of chaotropic agents (protein denaturants) including urea, guanidine hydrochloride, guanidine thiocyanate and formamide, which are incompatible with prior art indirect labeling systems wherein the structure of a biological macromolecule must be maintained to preserve requisite binding properties. Once formed, the bioconjugates are stable upon removal of water, and can be lyophilized for storage.

The stability of the bioconjugate at a given pH is determined to some extent by substituent group R_3 . Phenylboronic acid complexes of General Formula CVI, wherein group R_3 is H, are stable in buffered aqueous solutions over the approximate pH range 2.5 to 12.5. Phenylboronic acid complexes of General Formula CVI, wherein group R_3 is CH_3 , are stable in buffered aqueous solutions over the approximate pH range 3.5 to 12.5. Phenylboronic acid complexes of General Formula CVI, wherein group R_3 includes an electronegative moiety, are stable in buffered aqueous solutions over the approximate pH range less than 2.5 to 12.5.

The stability of the phenylboronic acid complex toward acid catalyzed hydrolysis is related to the pK_a of the hydroxamic acid participating in the


-96-

complex. The lower the pK_a of the hydroxamic acid moiety the more stable the complex. Consequently, phenylboronic acid complexes of General Formula CVI wherein group R_3 is an electronegative moiety exhibit greater stability toward acid catalyzed hydrolysis than do those in which R_3 is either H or CH_3 .

It should be appreciated that the pH stability associated with bioconjugates of General Formula CVI, derived from 4-aminomethyl-2,6-dihydroxybenzohydroxamic acid, is significantly better than that of bioconjugates of General Formula VI, derived from either 4- or 5-aminomethyl-salicylhydroxamic acid, with respect to both acid and base catalyzed hydrolysis.

The following examples present a detailed description of the synthesis of reagents of General Formula CI and General Formula CIII.

Example CI
Preparation of Methyl 4-Aminomethyl-2,6-
dihydroxybenzoate Hydrochloride



Methyl 4-Methyl-2,6-diacetoxybenzoate.

Methyl 4-methyl-2,6-dihydroxybenzoate (14.0 grams, 76.8 mmoles) was dissolved in anhydrous dichloromethane (250 mL) and acetic anhydride (21 mL, 223 mmoles) and anhydrous pyridine (18 mL, 223 mmoles) were added carefully. The solution was refluxed under dry nitrogen for 30 hours, then cooled to room temperature. The solution was washed twice with 1 M aqueous hydrochloric acid (200 mL portions) and then with saturated aqueous sodium chloride (200 mL). The dichloromethane solution was dried over anhydrous magnesium sulfate, filtered, and evaporated to an off-white solid. This crude product was flash

-97-

chromatographed on silica gel (250 grams) using hexanes:ethyl acetate (70:30 [v/v]) as the eluting solvent. Fractions containing the major product ($R_f = 0.3$) were pooled and evaporated to dryness to afford 19.0 grams (98% yield) of a white solid of methyl 4-methyl-2,6-diacetoxybenzoate (m.p. 70-71° C, open capillary, uncorrected).

^1H NMR (300 MHz, CHCl_3 -d) δ 2.27 (singlet, 6H, COCH_3), 2.37 (singlet, 3H, ArCH_3), 3.83 (singlet, 3H, OCH_3), 6.84 (singlet, 2H, ArH). ^{13}C NMR (75 MHz, CHCl_3 -d) δ 20.9, 21.5, 52.4, 116.5, 122.1, 143.8, 150.2, 163.9, 169.4.

Methyl 4-bromomethyl-2,6-diacetoxybenzoate.

Methyl 4-methyl-2,6-diacetoxybenzoate (14.9 grams, 56.0 mmoles) was dissolved in carbon tetrachloride (200 mL), and *N*-bromosuccinimide (11.1 grams, 62.2 mmoles) and benzoyl peroxide (0.2 grams, 0.8 mmoles) were added. The mixture was refluxed under nitrogen. After 3.5 hours, an additional portion (0.2 grams) of *N*-bromosuccinimide was added. Reflux was continued for an additional hour. The reaction mixture was cooled to room temperature and then in ice for 1 hour, and the solid removed by filtration. The filtrate was evaporated to dryness. Hexanes (250 mL) was added to the residue, and the mixture was boiled until dissolution was obtained. The hexanes solution was concentrated to about 75 mL when a solid just began to precipitate. The mixture was heated to dissolve the solid, and the solution was allowed to cool slowly to room temperature. White crystals formed slowly, and the mixture was chilled in ice for 2 hours to complete crystallization. The solid was filtered, washed with cold hexanes (100 mL), and dried *in vacuo* to afford 11.9 grams (62% yield) of methyl 4-bromomethyl-2,6-diacetoxybenzoate (m.p. 93-95° C, open capillary, uncorrected). The product was contaminated with a small amount of methyl 2,6-

-98-

diacetoxy-4-dibromomethylbenzoate, which did not interfere with subsequent reactions.

^1H NMR (300 MHz, CHCl_3-d) δ 2.29 (singlet, 6H, COCH_3), 3.85 (singlet, 3H, OCH_3), 4.41 (singlet, 2H, CH_2), 7.08 (singlet, 2H, ArH). ^{13}C NMR (75 MHz, CHCl_3-d) δ 20.9, 30.9, 52.7, 119.6, 121.8, 142.4, 150.2, 163.5, 169.1.

Methyl 4-aminomethyl-2,6-dihydroxybenzoate hydrochloride.

Methyl 4-bromomethyl-2,6-diacetoxybenzoate (45.8 grams, 133 mmoles) was dissolved in *N,N*-dimethylformamide (150 mL), and sodium azide (8.8 g, 135 mmoles) was added. The reaction mixture was stirred for 6 hours at room temperature. Dichloromethane (350 mL) was added to the reaction mixture, and this solution was washed with water (250 mL), 1 M aqueous hydrochloric acid (250 mL), and saturated aqueous sodium chloride (150 mL). The organic solution was dried over anhydrous magnesium sulfate, filtered, and evaporated to a clear, pale yellow oil. The oil was dissolved in methanol (750 mL) and transferred to a 2L Parr hydrogenation flask. Palladium on carbon catalyst (10% [w/w], 1.5 g) in water (10 mL) was added, followed by concentrated hydrochloric acid (15 mL). The flask was affixed to a Parr hydrogenator, and the mixture was shaken at room temperature under 30 psi of hydrogen for 24 hours. The reaction mixture was filtered through a 0.45 mm nylon filter. The retained solid was then washed with methanol (150 mL), and concentrated hydrochloric acid (7 mL) was added to the filtrate. The filtrate was heated to reflux for 2 hours, cooled to room temperature, and evaporated to dryness to give an off-white solid. This solid was dissolved in hot denatured ethanol (250 mL) and the solution was allowed to cool to room temperature. White crystals formed quickly. The mixture was then chilled for 16-18 hours at 4°C to complete crystallization. The solid was filtered, washed with a little


-99-

cold ethanol (50 mL) and then diethyl ether (150 mL), and dried *in vacuo* over potassium hydroxide pellets to afford 16.0 grams (52% yield based on monobromo starting material) of methyl 4-aminomethyl-2,6-dihydroxybenzoate hydrochloride (m.p. >260 °C, open capillary, uncorrected).

¹H NMR (300 MHz, DMSO-*d*₆) δ 3.77 (singlet, 3H, OCH₃), 3.83 (singlet, 2H, CH₂), 6.44 (singlet, 2H, ArH), 8.35 (broad singlet, 3H, NH₃), 10.20 (singlet, 2H, OH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 41.9, 52.1, 107.2, 107.3, 138.5, 157.6, 168.3.

Example CII

Preparation of a Reagent of General Formula CI Methyl (6-Aminohexanoyl)aminomethyl-2,6-dihydroxybenzoate Hydrochloride



Methyl (N-tert-butoxycarbonyl-6-aminohexanoyl)aminomethyl-2,6-dihydroxybenzoate.

Methyl 4-aminomethyl-2,6-dihydroxybenzoate hydrochloride (1.50 grams, 6.4 mmoles) was suspended in anhydrous *N,N*-dimethylformamide (25 mL), and *N,N*-diisopropylethylamine (2.2 mL, 12.8 mmoles) was added, followed by *N*-tert-butoxycarbonyl-6-aminohexanoic acid succinimidyl ester (2.10 grams, 6.4 mmoles). The mixture was stirred under dry nitrogen for 4 hours, during which time all solids dissolved. The solvent was then evaporated to leave a light brown syrup, which was partitioned between ethyl

-100-

acetate (70 mL) and 1 M aqueous hydrochloric acid (50 mL). The layers were separated, and the ethyl acetate solution was washed with saturated aqueous sodium bicarbonate (50 mL) and saturated aqueous sodium chloride (50 mL). The ethyl acetate solution was dried over anhydrous sodium sulfate, filtered, and evaporated to an amorphous off-white solid. The solid was crystallized from ethyl acetate/hexanes, filtered, and dried *in vacuo* to afford 2.10 grams (80% yield) of methyl (*N*-*tert*-butoxycarbonyl-6-aminohexanoyl)aminomethyl-2,6-dihydroxybenzoate (m.p. 73-76° C, open capillary, uncorrected).

¹H NMR (300 MHz, DMSO-*d*₆) δ 1.26 (multiplet, 2H, NHCH₂CH₂CH₂CH₂CH₂CO), 1.36 (multiplet, 2H, CH₂CH₂CO), 1.36 (singlet, 9 H, C(CH₃)₃), 1.50 (multiplet, 2 H, NHCH₂CH₂), 2.10 (triplet, *J* = 7 Hz, 2 H, CH₂CH₂CO), 2.87 (quartet, *J* = 6 Hz, 2 H, NHCH₂CH₂), 3.78 (singlet, 3 H, OCH₃), 4.08 (doublet, *J* = 6 Hz, 2 H, CH₂Ar), 6.23 (singlet, 2H, ArH), 6.75 (triplet, *J* = 6 Hz, 1 H, CONHCH₂CH₂), 8.26 (triplet, *J* = 6 Hz, 1 H, CONHCH₂Ar), 10.04 (singlet, 2H, OH). ¹³C NMR (75 MHz, CHCl₃-*d*) δ 25.0, 26.0, 28.3, 29.3, 35.3, 41.7, 52.0, 77.4, 104.3, 105.6, 145.8, 155.8, 158.2, 169.0, 172.4.

Methyl (6-aminohexanoyl)aminomethyl-2,6-dihydroxybenzoate hydrochloride.

Methyl (*N*-*tert*-butoxycarbonyl-6-aminohexanoyl)aminomethyl-2,6-dihydroxybenzoate (2.00 grmas, 4.87 mmoles) was dissolved in ethyl acetate (50 mL), and dry hydrogen chloride was bubbled slowly through the solution. The reaction mixture warmed as the gas dissolved. After 5 minutes, the gas was shut off, and the solution was stirred at room temperature. A white precipitate formed in the solution. After 30 minutes, the reaction mixture was chilled in ice for 2 hours, then the solid was filtered, washed with ethyl acetate (50 mL) and then diethyl ether (50 mL), and


-101-

dried *in vacuo* over potassium hydroxide pellets to afford 1.68 grams (99% yield) of methyl (6-aminohexanoyl)aminomethyl-2,6-dihydroxybenzoate hydrochloride (m.p. shrinks at 145° C, decomposes with effervescence at 152-154° C, open capillary, uncorrected).

¹H NMR (300 MHz, DMSO-*d*₆) δ 1.29 (multiplet, 2H, H₃CH₂CH₂CH₂CH₂CH₂CO), 1.55 (multiplet, 4H, NH₃CH₂CH₂CH₂CH₂CH₂CO), 2.13 (triplet, *J* = 8 Hz, 2H, CH₂CH₂CO), 2.70 (multiplet, 2H, NH₃CH₂CH₂), 3.75 (singlet, 3H, OCH₃), 4.07 (doublet, *J* = 6 Hz, 2H, CH₂Ar), 6.26 (singlet, 2H, ArH), 8.05 (broad singlet, 3H, NH₃), 8.39 (triplet, *J* = 6 Hz, 1H, CONHCH₂Ar), 9.84 (broad singlet, 2H, OH). ¹³C NMR (75 MHz, CHCl₃-*d*) δ 24.8, 25.6, 26.7, 35.1, 41.8, 52.0, 104.7, 105.5, 145.5, 158.1, 169.0, 172.3.

Example CIII

Preparation of a Reagent of General Formula CI Methyl *N*-Bromoacetyl-4-aminomethyl-2,6-dihydroxybenzoate



Methyl N-Bromoacetyl-4-aminomethyl-2,6-dihydroxybenzoate.

Bromoacetic acid (0.30 grams, 2.16 mmol) was dissolved in dry *N,N*-dimethylformamide (15 mL) and *N*-hydroxysuccinimide (0.25 grams, 2.21 mmol) and 1,3-dicyclohexylcarbodiimide (0.44 grams, 2.14 mmol) were added. The solution was stirred at room temperature for 1.25


-102-

hours. Methyl 4-aminomethyl-2,6-dihydroxybenzoate hydrochloride (0.50 grams, 2.14 mmol) was added, followed by *N,N*-diisopropylethylamine (0.38 mL, 2.15 mmol). The reaction mixture was stirred at room temperature for 2.5 hours, and was then diluted with ethyl acetate (100 mL). The mixture was filtered, and the filtrate washed twice with 1 N aqueous hydrochloric acid (100 mL then 50 mL) and with saturated aqueous sodium chloride (50 mL). The organic solution was dried over anhydrous magnesium sulfate, filtered, and evaporated to dryness. The crude product was purified by chromatography on silica gel, eluting with hexanes:ethyl acetate (1:1 [v/v]). Fractions containing pure product were pooled and evaporated to dryness to afford 0.43 grams of a white solid (63% yield) of methyl *N*-bromoacetyl-4-aminomethyl-2,6-dihydroxybenzoate (m.p. 157 - 158 °C).

¹H NMR (300 MHz, DMSO-*d*₆) δ 3.77 (singlet, 3H, OCH₃), 4.11 (singlet, 2H, BrCH₂), 4.13 (doublet, *J* = 6 Hz, 2H, CH₂NH), 6.25 (singlet, 2H, ArH), 8.70 (triplet, *J* = 6 Hz, 1H, CH₂NH), 10.05 (singlet, 2H, OH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 42.3, 42.6, 52.1, 104.7, 105.7, 144.7, 158.2, 166.3, 168.9.

Example CIV

Preparation of a Reagent of General Formula CI Methyl 4-Glutarylaminomethyl-2,6-dihydroxybenzoate Succinimidyl Ester



-103-

Methyl 4-glutarylaminomethyl-2,6-dihydroxybenzoate.

Glutaric anhydride (2.3 grams, 20 mmoles) was dissolved in dry tetrahydrofuran (100 mL), and the suspension was stirred under nitrogen. *N,N*-diisopropylethylamine (7.0 mL, 40 mmoles) was added, followed by methyl 4-aminomethyl-2,6-dihydroxybenzoate hydrochloride (4.7 grams, 20 mmoles). After stirring for 18 hours, the mixture was evaporated to dryness, and the resulting solid was dissolved in dichloromethane (100 mL) and allowed to sit at room temperature for 2 hours. The dichloromethane solution was evaporated to dryness to give a thick syrup, which was treated with cold 1 M aqueous hydrochloric acid (100 mL) for 30 minutes in ice. An off-white solid precipitated, which was filtered, washed with cold water, and dried *in vacuo* to afford 5.5 grams (88% yield) of methyl 4-glutarylaminomethyl-2,6-dihydroxybenzoate (m.p. 163-164° C).

¹H (300 MHz, DMSO-*d*₆) δ 1.74 (quintet, *J* = 8 Hz, 2H, CH₂CH₂CH₂), 2.16 (triplet, *J* = 8 Hz, 2H, CH₂CONH), 2.20 (triplet, *J* = 8 Hz, 2H, HO₂CCH₂), 3.78 (singlet, 3H, OCH₃), 4.10 (doublet, *J* = 6 Hz, 2H, ArCH₂NH), 6.23 (singlet, 2H, ArH), 8.31 (triplet, *J* = 6 Hz, 1H, NH), 10.05 (broad singlet, 3H, OH and CO₂H). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 20.7, 33.1, 34.4, 41.8, 52.1, 104.4, 105.6, 145.7, 158.2, 169.0, 172.0, 174.5.

Methyl 4-glutarylaminomethyl-2,6-dihydroxybenzoate succinimidyl ester.

Methyl 4-glutarylaminomethyl-2,6-dihydroxybenzoate (2.0 grams, 6.4 mmoles) was dissolved in dry tetrahydrofuran (80 mL), and *N*-hydroxysuccinimide (0.8 grams, 7.0 mmoles) and 1,3-dicyclohexylcarbodiimide (1.3 grams, 6.4 mmoles) were added. The mixture was stirred at room temperature under dry nitrogen, and the solids rapidly dissolved. After about 20 minutes, a white precipitate formed. The reaction mixture


-104-

was stirred 16-18 hours, then chilled several hours at -20°C . The mixture was filtered cold, and the solid washed with a little tetrahydrofuran (25 mL). The combined filtrates were evaporated to dryness, and the residue was crushed under ice-cold water (25 mL). The solid was filtered, washed quickly with cold water (25 mL) and then diethyl ether (100 mL), and dried *in vacuo*. The solid was recrystallized from ethyl acetate/hexanes to afford 2.5 grams (95% yield) of methyl 4-glutarylaminomethyl-2,6-dihydroxybenzoate succinimidyl ester (m.p. $161-163^{\circ}\text{C}$).

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 1.83 (quintet, $J = 8$ Hz, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 2.26 (triplet, $J = 8$ Hz, 2H, CH_2CONH), 2.73 (triplet, $J = 8$ Hz, 2H, NO_2CCH_2), 2.80 (singlet, 4H, $\text{COCH}_2\text{CH}_2\text{CO}$), 3.77 (singlet, 3H, OCH_3), 4.10 (doublet, $J = 6$ Hz, 2H, ArCH_2NH), 6.24 (singlet, 2H, ArH), 8.35 (triplet, $J = 6$ Hz, 1H, NH), 10.04 (broad singlet, 2H, OH). ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 20.4, 25.4, 29.7, 33.6, 41.8, 52.0, 104.5, 105.6, 145.5, 158.1, 168.9, 170.5, 171.4.

Example CV

Preparation of a Reagent of General Formula C Methyl 4-Glutarylaminomethyl-2,6-dihydroxybenzoic Acid Hydrazide



-105-

Methyl 4-glutarylaminomethyl-2,6-dihydroxybenzoate N-tert-butyloxycarbonylhydrazide.

Methyl 4-glutarylaminomethyl-2,6-dihydroxybenzoate succinimidyl ester (1.6 grams, 3.9 mmoles) was suspended in dry dichloromethane (50 mL), and *tert*-butylcarbazate (0.50 grams, 3.9 mmoles) was added. The reaction was stirred at room temperature for 24 hours. The pale solution was then washed twice with saturated aqueous sodium bicarbonate (50 mL portions), 0.1 M aqueous hydrochloric acid (50 mL), and saturated aqueous sodium chloride (50 mL). It was then dried over anhydrous sodium sulfate, filtered, and evaporated to dryness. The residue was flash chromatographed on a 2.5 X 150 cm column of silica gel, eluting with ethyl acetate. Fractions containing pure product were pooled and evaporated to a clear syrup. This syrup was crystallized from ethyl acetate:hexanes at -20° C. The solid was filtered, washed with cold hexanes, and dried *in vacuo* to afford 1.0 grams (62% yield) of methyl 4-glutarylaminomethyl-2,6-dihydroxybenzoate *N-tert*-butyloxycarbonylhydrazide (m.p. 95-100° C, softens at 88° C, open capillary, uncorrected).

¹H (300 MHz, DMSO-*d*₆) δ 1.37 (singlet, 9H, (CH₃)₃C), 1.74 (quintet, *J* = 7 Hz, 2H, CH₂CH₂CH₂), 2.07 (triplet, *J* = 7 Hz, 2H, CH₂CONHCH₂), 2.15 (triplet, *J* = 7 Hz, 2H, HNHNOCCCH₂), 3.77 (singlet, 3H, OCH₃), 4.09 (doublet, *J* = 6 Hz, 2H, ArCH₂NH), 6.23 (singlet, 2H, ArH), 8.28 (triplet, *J* = 6 Hz, 1H, NH), 8.65 (singlet, 1H, NHNHCOCH₂), 9.47 (singlet, OCONHNH), 9.96 (singlet, 2H, OH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 21.2, 28.0, 32.7, 34.7, 41.7, 52.1, 79.1, 104.4, 105.6, 145.7, 155.5, 158.2, 169.0, 171.7, 172.0.

Methyl 4-Glutarylaminomethyl-2,6-dihydroxybenzoic Acid Hydrazide.

Methyl 4-glutarylaminomethyl-2,6-dihydroxybenzoate *N-tert*-butyloxycarbonyl hydrazide (0.50 grams, 1.2 mmoles) is


-106-

dissolved in dry tetrahydrofuran (25 mL), and dry hydrogen chloride was bubbled slowly through the solution. The reaction mixture warmed as the gas dissolved. After 5 minutes, the gas was shut off, and the solution was stirred at room temperature. A white precipitate formed in the solution. After 30 minutes, the reaction mixture was chilled in ice for 2 hours, then the solid was filtered, washed with diethyl ether (50 mL), and dried *in vacuo* over potassium hydroxide pellets to afford 0.42 grams (98% yield) of methyl 4-glutarylaminomethyl-2,6-dihydroxybenzoic acid hydrazide (m.p. 158-159° C, decomposed with effervescence, open capillary, uncorrected).

^1H (300 MHz, $\text{DMSO}-d_6$) δ 1.76 (quintet, $J = 7$ Hz, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 2.16 (triplet, $J = 7$ Hz, 2H, $\text{CH}_2\text{CONHCH}_2$), 2.21 (triplet, $J = 7$ Hz, 2H, $\text{H}_3\text{NHNOCCH}_2$), 3.75 (singlet, 3H, OCH_3), 4.07 (doublet, $J = 6$ Hz, 2H, ArCH_2NH), 5.00-7.00 (very broad singlet, 2H, OH), 6.25 (singlet, 2H, ArH), 8.39 (triplet, $J = 6$ Hz, 1H, NH), 10.49 (broad singlet, 3H, NHNH_3), 11.09 (singlet, 1H, NHNH_3). ^{13}C NMR(75 MHz, CHCl_3-d) δ 20.9, 32.2, 34.4, 41.8, 52.0, 104.8, 105.6, 145.5, 158.1, 169.0, 171.6, 171.9.

Example CVI

Preparation of a Reagent of General Formula CIII Synthesis of 4-Glutarylaminomethyl-2,6-dihydroxybenzohydroxamic Acid Hydrazide



-107-

*4-Glutarylaminomethyl-2,6-dihydroxybenzohydroxamic Acid
N-tert-Butyloxycarbonylhydrazide.*

Methyl 4-glutarylaminomethyl-2,6-dihydroxybenzoate *N-tert*-butyloxycarbonylhydrazide (0.30 grams, 0.7 mmoles) was added to a cooled (ice/water bath) solution of hydroxylamine sulfate (0.12 grams, 0.7 mmoles), sodium hydroxide (0.15 grams, 3.6 mmoles) and sodium sulfite (0.05 grams, 0.4 mmoles) in water (6 mL). The suspension was stirred for 24 hours in the dark, allowing it to warm to room temperature. The solution was dilute with 1 M hydrochloric acid (4 mL), and extracted twice with ethyl acetate (15 mL portions). The combined ethyl acetate extracts were washed with water (10 mL) and saturated aqueous sodium chloride (10 mL), dried over anhydrous magnesium sulfate, filtered, and evaporated to dryness. The residue was dissolved in a minimum volume of ethyl acetate, and diethyl ether was added to precipitate an orange gummy material. The gum was triturated under dry diethyl ether to produce a solid, which was filtered, washed with ether, and dried *in vacuo* to afford 0.21 grams (69% yield) of 4-glutarylaminomethyl-2,6-dihydroxybenzohydroxamic acid *N-tert*-butyloxycarbonylhydrazide (m.p. decomposed with effervescence at 60-65° C, open capillary, uncorrected).

¹H (300 MHz, DMSO-*d*₆) δ 1.37 (singlet, 9H, (CH₃)₃C), 1.73 (quintet, *J* = 7 Hz, 2H, CH₂CH₂CH₂), 2.07 (triplet, *J* = 7 Hz, 2H, CH₂CONHCH₂), 2.15 (triplet, *J* = 7 Hz, 2H, HNHNOCCCH₂), 4.08 (doublet, *J* = 6 Hz, 2H, ArCH₂NH), 6.22 (singlet, 2H, ArH), 8.28 (triplet, *J* = 6 Hz, 1H, NH), 8.65 (singlet, 1H, NHNHCOCH₂), 9.47 (singlet, OCONHNH), 10.77 (broad singlet, 1H, NHOH), 12.30 (broad singlet, 3H, ArOH and NHOH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 21.3, 28.1, 32.7, 34.7, 41.7, 79.1, 100.1, 105.6, 145.8, 155.6, 159.8, 167.5, 171.7, 172.0.

-108-

4-Glutarylaminomethyl-2,6-dihydroxybenzohydroxamic acid hydrazide trifluoroacetate.

4-Glutarylaminomethyl-2,6-dihydroxybenzohydroxamic acid *tert*-butyl carbazate (0.09 grams, 0.2 mmoles) was suspended in dry dichloromethane (5 mL), and trifluoroacetic acid (0.5 mL) was added. The solid dissolved on addition of the acid, and the reaction was stirred at room temperature for three hours. After one hour, an orange precipitate had formed. The precipitate was collected by filtration and dried *in vacuo* over potassium hydroxide pellets to afford 0.10 grams (100% yield) of 4-glutarylaminomethyl-2,6-dihydroxybenzohydroxamic acid hydrazide trifluoroacetate (m.p. 104-108° C, open capillary, uncorrected).

¹H (300 MHz, DMSO-*d*₆) δ 1.80 (quintet, *J* = 7 Hz, 2H, CH₂CH₂CH₂), 2.17 (triplet, *J* = 7 Hz, 2H, CH₂CONHCH₂), 2.23 (triplet, *J* = 7 Hz, 2H, HNHNOCCCH₂), 4.09 (doublet, *J* = 6 Hz, 2H, ArCH₂NH), 6.22 (singlet, 2H, ArH), 8.32 (triplet, *J* = 6 Hz, 1H, NH), 8.95 - 10.60 (very broad singlet, 5H, NHHNH₃ and NHOH), 11.80-12.95 (very broad singlet, 3H, ArOH and NHOH). ¹³C NMR (75 MHz, DMSO-*d*₆) δ 20.9, 32.2, 34.4, 41.8, 100.1, 105.7, 145.8, 158.5, 159.0, 159.9, 167.5, 171.8, 171.9.

Example CVII

Preparation of Conjugates of General Formula CIV

Synthesis of 5'-PBA-labeled Oligodeoxyribonucleotide Conjugates.

Oligodeoxyribonucleotide 7172 (sequence 5'-GATTACGCCAGTTGTACGGAC-3') was synthesized on a 1 μmole scale using standard automated phosphoramidite chemistry (Beckman Instruments Oligo 1000 and associated reagents). A protected amine-containing phosphor-amidite (Aminolink 2, Applied Biosystems or UniLink Amino Modifier, Clontech) was

-109-

employed on the same instrument to introduce one to four, reactive primary amines onto the 5'-end of the oligodeoxyribonucleotide using standard chemistry. The completed oligodeoxyribo-nucleotide was then cleaved from the support and the nucleobases deprotected using an UltraFast Deprotection kit (Beckman Instruments) and the protocol supplied by the manufacturer.

The amino-oligonucleotides were purified by ethanol precipitation, dissolved in 0.8 mL of 0.1 M NaHCO_3 , and condensed with phenylboronic acid reagent (PBA-Z-NHS) having a reactive *N*-hydroxysuccinimidyl ester moiety (5 mgs per mmole of primary amino groups on the amino-oligonucleotide in 0.2 mL of anhydrous *N,N*-dimethylformamide) for 2-18 hours at room temperature.

The crude PBA-modified oligonucleotide was isolated from the reaction mixture by gel filtration on a KwikSep Dextran column (Pierce Chemical) in 0.1 M aqueous triethylammonium acetate, pH 6.5. The PBA-modified oligonucleotide was then concentrated in a vacuum centrifuge to 1 mL, and purified by reverse phase HPLC on a 4.6 mm x 250 mm C18 column, with a triethylammonium acetate-acetonitrile gradient. The desired product peak was collected and evaporated to a small pellet in a vacuum centrifuge, dissolved in 0.5 mL of water, and stored frozen.

Preparation of Salicylhydroxamic Acid Magnetic Beads.

Ten milliliters of unmodified M280 or M450 magnetic beads (Dyna) were gradually dehydrated into acetonitrile, and converted to aldehyde modified beads using oxalyl chloride activated *N,N*-dimethylsulfoxide and triethylamine in dichloromethane at -78°C . The resulting aldehyde bearing beads were gradually rehydrated and suspended in 5 mL of 0.1 M sodium acetate pH 5.5. The aldehyde groups were coupled with 4-glutarylaminomethylsalicyl-hydroxamic acid

-110-

hydrazide (SHA-Z-NHNH₂) by adding 15-25 mgs dissolved in 200 μ L *N,N*-dimethylsulfoxide, and rotating coupling reaction over night at room temperature. The beads were then washed extensively with water and stored in 5 mL of 10% ethanol.

Preparation of Salicylhydroxamic Acid (SHA) Sepharose 4B.

SHA-Sepharose 4B was prepared by mixing 130 mg of (6-aminohexanoyl)-4-amino-methylsalicylhydroxamic acid (SHA-Z-NH₂), dissolved in 30 mL 0.2 M NaHCO₃, with 6.5 g HCl washed CNBr activated Sepharose 4B (Pharmacia) overnight at room temperature. After the coupling reaction, 2 mL of 0.5 M Tris, pH 8.5 were added and the gel slurry mixed at room temperature for 1 hour, and washed with water, 0.5 M NaCl, and water again. The resulting SHA-Sepharose 4B was suspended in 30 mL of 20% ethanol, and stored at 4° C.

Preparation of 2,6-Dihydroxybenzohydroxamic Acid (DHBHA) Sepharose 4B.

DHBHA-Sepharose 4B was prepared by mixing 114 mg methyl (6-amino-hexanoyl)-4-aminomethyl-2,6-dihydroxybenzoate [DHBA(OMe)-X-NH₂], dissolved in 30 mL 0.2 M NaHCO₃, with 5 g HCl washed CNBr activated Sepharose 4B (Pharmacia), overnight at room temperature. After the coupling reaction, the gel was washed with water and suspended in 50 mL 0.1 M NH₂OH, pH 9, and rotated at room temperature for two hours. Finally, the gel was washed with water and suspended in 30 mL of 20% ethanol, and stored at 4°C.

Preparation of a Phenylboronic Acid- α -Biotin Antibody Conjugate.

One milliliter of α -Biotin monoclonal IgG₁ antibody (6.5 mg/mL in 0.1 M NaHCO₃) was conjugated with 440 nmoles of PBA-Z-NHS (7.4 μ L of 60 mM PBA-Z-NHS dissolved in

-111-

N,N-dimethylsulfoxide) for 1 hour at room temperature. Unconjugated PBA-Z-NHS and its hydrolysis products were removed by dialysis. The ultra-violet absorbance spectrum of the resulting conjugate (PBA- α -Biotin) exhibited an increase in A_{260} relative to A_{280} consistent with phenylboronic acid modification.

Preparation of a Phenylboronic Acid-Alkaline Phosphatase Conjugate.

One milliliter of alkaline phosphatase (Sigma, 6 mg/mL) was dialyzed against one liter of 0.1 M NaHCO_3 , and conjugated with 700 nmoles of PBA-Z-NHS (10 μL of 70 mM stock in *N,N*-dimethylformamide) for two hours on ice. Unconjugated PBA-Z-NHS and its hydrolysis products were removed by dialysis in 0.1 M NaHCO_3 . The ultra-violet absorbance spectrum of the resulting conjugate (PBA-AP) exhibited an increase in A_{260} relative to A_{280} consistent with phenylboronic acid modification. The conjugate was stored at 4° C.

Preparation of a 2,6-Dihydroxybenzohydroxamic Acid-Alkaline Phosphatase Conjugate.

One milliliter of alkaline phosphatase (Sigma, 6 mg/mL) was dialyzed against one liter of 0.1 M NaHCO_3 , and conjugated with 714 nmoles of methyl 4-glutarylaminomethyl-2,6-dihydroxybenzoate succinimidyl ester [DHBA(OMe)-Z-NHS] (10.5 μL of 68 mM in *N,N*-dimethylformamide) for two hours on ice. The methyl ester of the conjugate was converted to a hydroxamic acid by adding one milliliter of 2 M NH_2OH , pH 10, and incubating the reaction at 4° C for six days. The NH_2OH reaction mixture was then dialyzed against 0.1 M NaHCO_3 and stored at 4° C.

-112-

Preparation of a Salicylhydroxamic Acid-Goat α -Mouse Antibody Conjugate.

Two milliliters of goat α -mouse antibody (Rockland, 8.8 mg/mL in 0.1 M NaHCO_3) were conjugated with 2.35 umoles of methyl 4-glutarylaminomethylsalicylate succinimidyl ester [SA(OMe)-Z-NHS] for 1 hour at room temperature. The methyl ester of the conjugate was converted to a hydroxamic acid by adding two milliliters of 2 M NH_2OH , pH 10, adjusting the pH to 10 with 1 N NaOH, and incubating the reaction at room temperature for three days. NH_2OH and unconjugated SA(OMe)-X-NHS and its hydrolysis products were removed by gel filtration on a G-25 Sephadex column (Pharmacia) in 0.1 M NaHCO_3 , and the conjugate (SHA-goat α -mouse) was stored at 4° C.

Preparation of a 2,6-Dihydroxybenzohydroxamic Acid-Goat α -Mouse Antibody.

Two milliliters of goat α -mouse antibody (Rockland, 8.8 mg/mL in 0.1 M NaHCO_3) were conjugated with 2.35 umoles of DHBA(OMe)-Z-NHS for 1 hour at room temperature. The methyl ester of the conjugate was converted to a hydroxamic acid by adding two milliliters of 2 M NH_2OH , pH 10, adjusting the pH to 10 with 1 N NaOH, and incubating the reaction at room temperature for three days. NH_2OH and unconjugated DHBA(OMe)-X-NHS and its hydrolysis products were removed by gel filtration on a G-25 Sephadex column (Pharmacia) in 0.1 M NaHCO_3 , and the conjugate (DHBHA-goat α -mouse) stored at 4° C.

Polymerase Chain Reaction (PCR) Protocol.

A region of Lambda DNA (801bp) was amplified by the polymerase chain reaction. The PCR reaction contained 200 μM dATP, dCTP, dGTP, and dTTP in addition to Biotin- and PBA- modified oligonucleotide primers at 1 μM in 1X PCR

-113-

Buffer II (Perkin Elmer), Lambda DNA (1 ng/uL), and 1U of *Thermus aquaticus* DNA polymerase. The reaction mixture was denatured at 92° C for one minute and amplified by 35 cycles of PCR at 95° C for 10 seconds, 62° C for 20 seconds, and 72° C for 30 seconds, with a final extension at 72° C for 5 minutes. The reaction produced 50-100 ng of amplified product (801bp), which exhibited retarded mobility relative to unmodified PCR product during electrophoresis on 1% agarose gels in 50 mM Tris, 100 mM borate, 2 mM EDTA, pH 8.3.

Example CVIII

Preparation of Bioconjugates of General Formula CVI Sandwich Hybridization Detection of Nucleic Acid Probes

A 42-mer oligonucleotide was hybridized with two 21-mer oligonucleotides bearing 5'-PBA and Biotin labels in 1.5 M NaCl, 150 mM sodium citrate, pH 7, at 45° C for ten minutes. Twenty-five microliters of the hybridization mixture was mixed with twenty five microliters of M280 streptavidin-magnetic particles (Dyna) in a polypropylene microtiter plate well. After 30 minutes, the magnetic particles were captured in the bottom of the well with a magnetic plate, and washed five times with 150 mM NaCl, 20 mM Tris-HCl, 0.02% Tween20, pH 8.

One hundred microliters of DHBHA-AP (1 ug/mL in 1 mg/mL BSA, 140 mM NaCl, 10 mM Tris-HCl, pH 8) were added to the magnetic particles and mixed well. After 30 minutes, the magnetic particles were captured in the bottom of the well with a magnetic plate, and washed six times with 150 mM NaCl, 20 mM Tris-HCl, 0.02% Tween20, pH 8. Alkaline phosphatase substrate (1 mg/mL *p*-nitrophenylphosphate in 1 M diethanolamine buffer, 1 mM MgCl₂, 0.1 mM ZnCl₂, pH 10.4)

-114-

was added, and incubated at 37° C for 90 minutes. The Absorbance at 405 nm (A_{405}) was measured with an ELISA plate reader (Molecular Devices).

A strong A_{405} was produced when all components of the hybridization sandwich were present, and the signal was proportional to the amount of 42-mer present. Experimental controls lacking either the 42-mer, the Biotin-oligonucleotides and PBA-oligonucleotides did not produce a significant A_{405} .

Example CIX

Preparation of Bioconjugates of General Formula CVI

Detection of PBA-Labeled PCR Product on DHBHA-Coated Microtiter Plates.

The wells of a polystyrene microtiter plate (Becton Dickinson) were coated with DHBHA by filling the wells with 200 μ L of DHBHA-goat α -mouse conjugate (30 μ g/mL in 0.1M NaHCO_3 pH 9.0) and incubating overnight at 4° C. The coating solution was removed and the plate backcoated with 5 mg/mL BSA (300 μ L per well in 0.2 M NaHCO_3 , pH 9.0) for 1 hour at room temperature. The BSA solution was removed by washing the plate five times with ELISA Wash Buffer (150 mM NaCl, 20 mM Tris-HCl, .02% Tween 20, pH 8.0).

One hundred microliters of unpurified PBA and biotin labeled PCR product were added to 900 μ L of 1.5N NaCl, 150 mM sodium citrate, pH 7.0 (10X SSC) and serially-diluted in 10X SSC. One hundred microliters of the diluted PCR products were added to the wells and incubated for one hour at room temperature. The plate was then washed five times with ELISA Wash Buffer, and 100 μ L of Streptavidin-Alkaline Phosphatase (Boehringer Mannheim, 0.2 U/mL in 1 mg/mL BSA,

-115-

140 mM NaCl, 10 mM Tris-HCl, pH 8.0) were added to each well and incubated for thirty minutes at room temperature.

The plate was washed 5 times with ELISA Wash Buffer, and 200 μ L of *p*-nitrophenyl- phosphate (1 mg/mL in diethanolamine, 1 mM $MgCl_2$, 0.1 mM $ZnCl_2$, pH 10.4) were added to the plate and incubated at 37° C for 30-60 minutes. Less than 1 μ L of PCR product was detected. PCR product lacking either PBA or biotin labels was not detected.

Example CX

Preparation of Bioconjugates of General Formula CVI Detection of PBA-Labeled PCR Product

Detection of PBA-Labeled PCR Product on SHA-Magnetic Beads.

PBA-labeled PCR product (0.02 μ L - 5 μ L) was diluted into 25-100 μ L of 1.5 M NaCl, 150 mM sodium citrate, pH 7 (10X SSC), and added to a polypropylene microtiter plate well containing SHA-magnetic particles (10-50 μ L). The particles and PCR product were mixed occasionally for 30-60 minutes at room temperature. The magnetic particles were captured in the bottom of the wells with a magnetic plate and washed five times in 150 mM NaCl, 20 mM Tris-HCl, 0.02% Tween 20, pH 8 (ELISA Wash Buffer). One hundred microliters of streptavidin alkaline-phosphatase (Boehringer Mannheim, 0.2 U/mL in 1 mg/mL BSA, NaCl, Tris-HCl, pH 8) were added and mixed with the magnetic particles for 30 minutes at room temperature. The magnetic particles were captured in the bottom of the wells with a magnetic plate and washed 5 times with ELISA Wash. Alkaline phosphatase substrate was added (1 mg/mL *p*-nitrophenyl phosphate in 1M diethanolamine buffer, 1 mM $MgCl_2$, 0.1 mM $ZnCl_2$, pH 10.4), and the color

-116-

developed at 37° C for 10-60 minutes. The lower limit of detection was 50 pg of PCR product.

Example CXI

Preparation of Bioconjugates of General Formula CVI Detection of a PBA-Labeled Oligonucleotide Hybrid

Detection of a PBA-Labeled Oligonucleotide Hybridized to a Biotin-Labeled Oligonucleotide.

A 42-mer oligonucleotide was hybridized with two 21-mer oligonucleotides bearing 5'-PBA and Biotin labels in 1.5 M NaCl, 150 mM sodium citrate, pH 7, at 45 C for ten minutes. Twenty-five microliters of the hybridization mixture was mixed with 1-50 uL of SHA-magnetic particles (DynaL, M450) in a polypropylene microtiter plate well. After 30 minutes, the magnetic particles were captured in the bottom of the well with a magnetic plate, and washed five times with 150 mM NaCl, 20 mM Tris-HCl, 0.02% Tween 20, pH 8.

One hundred microliters of SHA-AP (1 ug/mL in 1 mg/mL BSA, 140 mM NaCl, 10 mM Tris-HCl, pH 8) were added to the magnetic particles and mixed well. After 30 minutes, the magnetic particles were captured in the bottom of the well with a magnetic plate, and washed six times with 150 mM NaCl, 20 mM Tris-HCl, 0.02% Tween 20, pH 8. The particles were mixed with alkaline phosphatase substrate (1 mg/mL *p*-nitrophenyl phosphate in 1 M diethanolamine buffer, 1 mM MgCl₂, 0.1 mM ZnCl₂, pH 10.4) and incubated at 37° C for 90 minutes. The A₄₀₅ was measured with a ELISA plate reader (Molecular Devices). As little as 45 pg of oligonucleotide 42-mer was detected. Experimental controls lacking either the 42-mer, or the PBA or Biotin labeled oligonucleotides did not produce a significant A₄₀₅.

-117-

Example CXII**Preparation of Bloconjugates of General Formula CVI*****Immobilization of a PBA- α -Biotin Conjugate on SHA-Sepharose 4B.***

One mg of PBA- α -Biotin, diluted to 1 mL with Tris buffered saline, was applied to small column of SHA-Sepharose 4B (1.0 x 2.0 cm), and washed extensively with Tris buffered saline. The size of the A_{280} peak of the material not binding to the column indicated that almost all of the PBA-conjugate was immobilized on the column.

Biotin binding activity of the column was assayed by applying to the column 5 mL of 1 ug/mL biotinylated alkaline phosphatase in Tris buffered saline containing 5 mg/mL bovine serum albumin (BSA). A sample of the peak of the material flowing through the column was collected for comparison of the enzymatic activity with a sample of the alkaline phosphatase dilution applied to column. After applying the sample, the column was washed with 20 mL of buffer. After washing, a very small sample of column material (25 uL liquid containing about 1 uL gel) was collected to measure the enzymatic activity bound to the gel as a result of capture by the immobilized α -biotin antibody.

The alkaline phosphatase activity was measured by incubating 25 uL of the enzyme samples in 250 uL of 1 mg/mL p-nitrophenylphosphate in 1 M diethanolamine buffer, 1 mM $MgCl_2$, and 0.1 mM $ZnCl_2$, pH 10.4, for 20 minutes and then adding 650 uL of 0.1 M $NaHCO_3$, 10 mM EDTA. Relative to a buffer blank, the A_{405} of the sample applied to the column was 1.57, while the A_{405} of the peak of the material not retained by the column was only 0.042, indicating that virtually all the enzyme conjugate was captured by the column. The small

-118-

amount of gel assayed produced an A_{405} of 1.30, demonstrating that the enzyme was in fact captured by the column.

Example CXIII

Preparation of Bioconjugates of General Formula CVI


Immobilization of a PBA-Alkaline Phosphatase Conjugate on SHA-Magnetic Beads.

PBA-conjugated alkaline phosphatase was diluted to 5 ug/mL in Tris buffered saline containing 5 mg/mL bovine serum albumin. Two hundred microliters of diluted PBA-conjugated enzyme were mixed with 5, 10, or 20 uL of SHA-magnetic beads (Dyna, M280). The enzyme was also mixed with 40 uL of unmodified beads as a control. The beads were mixed gently for 10 minutes on ice, after which the beads were captured with a rare earth magnet and washed 4 times with Tris buffered saline. The beads were then suspended in 250 uL of 1 mg/mL *p*-nitrophenylphosphate in 1 M diethanolamine buffer, 1 mM $MgCl_2$, and 0.1 mM $ZnCl_2$, pH 10.4, and mixed occasionally at 37° C for 10 minutes. The reactions were terminated with 750 uL of Tris buffered saline, 5 mM EDTA. The A_{405} relative to a buffer blank was measured to determine the alkaline phosphatase activity bound to the magnetic beads. The control beads produced an A_{405} of only 0.15, while the SHA-magnetic beads produced an A_{405} of 0.62, 0.97, and 1.33 for 5, 10, and 20 uL of beads, respectively, indicating the immobilization of significant amounts of PBA-AP conjugate on the surface of the beads.

-119-

IN THE CLAIMS

1. A reagent having the general formula of General Formula CI:




General Formula CI

wherein group Z comprises a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide and disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length;

wherein group R₁ is a electrophilic or nucleophilic moiety suitable for reaction of the reagent with a bioactive species; and

wherein group R₂ is one of an alkyl and a methylene group with an electronegative moiety.


2. The reagent of claim 1, wherein group R₂ is preferably selected from one of CH₃, CH₂CH₃, CH₂CN, CH₂COOH, CH₂CONH₂, and CH₂OCH₃.
3. The reagent of claim 1, wherein group Z is preferably an unbranched alkyl chain of the general formula (CH₂)_n, wherein n = 1 to 6.
4. The reagent having the formula:



wherein group R₂ is one of an alkyl and a methylene with an electronegative moiety.

-120-

5. The reagent of claim 1, wherein group R_1 is preferably selected from the group consisting of acrylamide, amino, bromo, dithiopyridyl, bromoacetamide, hydrazide, N-hydroxysuccinimide ester, N-hydroxysulfosuccinimide ester, imidate ester, imidazolide, iodo, iodoacetamide, maleimide, and thiol moieties.
6. A conjugate of a biologically active species with a reagent having the general formula of General Formula CII:



General Formula CII

wherein group Z comprises a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide and disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length;


wherein group R_2 is one of an alkyl and a methylene with an electronegative moiety; and

wherein BAS is a biologically active species.

7. The conjugate of claim 6, wherein group R_2 is preferably selected from one of CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 , and CH_2OCH_3 .
8. The conjugate of claim 6, wherein group Z is preferably an unbranched alkyl chain of general formula $(\text{CH}_2)_n$, wherein $n = 1$ to 6.


-121-

9. The conjugate of claim 6, having the general formula:



wherein group R_2 is CH_3 .


10. A compound comprising:



wherein R_2 is one of an alkyl and a methylene with an electronegative moiety.

11. The compound of claim 10, wherein group R_2 is preferably selected from one of CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 , and CH_2OCH_3 .


12. A method for preparing a reagent, the method comprising:
condensing methyl 2,6-dihydroxy-4-methylbenzoate with a suitable protecting groups Q to modify the reactivity of the ring system:




wherein R_2 is an alkyl,

-122-


halogenating the alkyl benzoate ester to yield a benzyl halide wherein group Y is a halogen:



alkylating the benzyl halide with an azide salt to yield a benzyl azide:




and hydrogenating the benzyl azide to yield the corresponding benzyl amine and remove the protective groups Q:



13. The method of claim 12, wherein the protective group Q is an acetyl group.
14. The method of claim 12, further comprising:
condensing the 2,6-dihydroxybenzylamine with one of an activated carboxylic acid to yield the reagent of general formula:

-123-




wherein group Z comprises a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide and disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length; and

wherein group R₁ is a electrophilic or nucleophilic moiety suitable for reaction of the reagent with a bioactive species.

15. The method of claim 12, wherein group R₁ is preferably selected from the group consisting of acrylamide, amino, bromo, dithiopyridyl, bromoacetamide, hydrazide, N-hydroxysuccinimide ester, N-hydroxysulfosuccinimide ester, imidate ester, imidazolidine, iodo, iodoacetamide, maleimide, and thiol moieties.


16. The method of claim 14, further comprising the step of condensing the reagent with a bioactive species to yield the reagent of general formula:



wherein BAS is a biologically active species.

-124-

17. A method for preparing the reagent of General Formula CI:




General Formula CI

wherein group Z comprises a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with one of intermediate amide and disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length;

wherein group R_1 is a electrophilic or nucleophilic moiety suitable for reaction of the reagent with a bioactive species and

wherein group R_2 is is one of an alkyl and a methylene with an electronegative moiety,
the method comprising:


condensing a 2,6-dihydroxy-4-methylbenzoate with a suitable protective group Q to modify the reactivity of the ring system:




wherein group X is an alkyl group of from about 1 to 3 carbons,

halogenating the 2,6-dihydroxy-4-methylbenzoate to yield a benzyl halide
wherein group Y is a halogen:


-125-




alkylating the benzyl halide with an azide salt to yield a benzyl azide:



hydrogenating the benzyl azide and removing the protecting groups Q to yield a benzyl amine:




condensing the benzyl amine with a suitable protective group Q' to protect the amine moiety from participation in subsequent reactions, to yield an alkyl aminomethylbenzoate:




hydrolyzing the alkyl ester and to yield a benzoic acid:

-126-




alkylating the benzoic acid to yield an ester:



wherein group R_2 is selected from one of an alkyl and a methylene with an electronegative moiety,

removing the protective group Q' to yield a benzyl amine:



condensing the benzylamine with one of an activated carboxylic acid to yield the reagent.


18. The method of claim 17, wherein group R_1 is preferably selected from the group consisting of acrylamide, amino, bromo, dithiopyridyl, bromoacetamide, hydrazide, N-hydroxysuccinimide ester, N-hydroxysulfosuccinimide ester, imidate ester, imidazolidine, iodo, iodoacetamide, maleimide, and thiol moieties.

19. The method of claim 17, wherein group R_2 is selected from one of CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 , and CH_2OCH_3 .

20. The method of claim 17, wherein the protective group Q is an acetyl group.


-127-

21. The method of claim 17, wherein the protective group Q' is a N-(tert-butoxycarbonyl) group.
22. The method comprising the step of condensing the reagent of General Formula XI as in claim 1 with a bioactive species to yield the reagent of General Formula XII:



wherein *BAS* is a biologically active species.

23. A reagent having the general formula of General Formula CIII:



General Formula CIII

wherein group Z comprises a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide and disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length;

wherein group R₃ is selected from one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent; and


wherein group R₁ is an electrophilic or nucleophilic moiety suitable for reaction of the reagent with a bioactive species.

-128-

24. The reagent of claim 23, wherein group R_3 is preferably selected from one of H, CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 , $\text{CH}_2\text{CH}_2\text{OH}$, and CH_2OCH_3 .

25. The reagent of claim 23, wherein group Z is preferably an unbranched alkyl chain of the general formula $(\text{CH}_2)_n$, wherein $n = 1$ to 6.


26. The reagent having the general formula:



wherein group R_3 is one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent.

27. The reagent of claim 23, wherein group R_1 is preferably selected from the group consisting of acrylamide, amino, bromo, dithiopyridyl, bromoacetamide, hydrazide, N-hydroxysuccinimide ester, N-hydroxysulfosuccinimide ester, imidate ester, imidazolidine, iodo, iodoacetamide, maleimide, and thiol moieties.

28. A conjugate of a biologically active species with a reagent having the general formula of General Formula CIV:



General Formula CIV

wherein group Z comprises a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length


-129-

with at least one of intermediate amide and disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length; wherein group R_3 is selected from one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent; and wherein *BAS* is a biologically active species.

29. The conjugate of claim 28, wherein group R_3 is preferably selected from one of H, CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 , CH_2CH_2OH , and CH_2OCH_3 .


30. The conjugate of claim 28, wherein group *Z* is preferably an unbranched alkyl chain of general formula $(CH_2)_n$, wherein $n = 1$ to 6.

31. The conjugate of claim 28, having the general formula:



wherein group R_3 is H.

32. A bioconjugate having the general formula of General Formula CVI:



General Formula CVI

wherein group *Z* comprises a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched

-130-

saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with one of intermediate amide and disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length;

wherein group R_3 is selected is one of H, an alkyl, and a methylene or ethylene with an electronegative moiety;

wherein *BAS* is a biologically active species; and


wherein *BAS** is a biologically active species.

33. The bioconjugate of claim 32, wherein group R_3 is preferably selected from one of H, CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 , CH_2CH_2OH , and CH_2OCH_3 .

34. The bioconjugate of claim 32, wherein group Z is preferably an unbranched alkyl chain of general formula $(CH_2)_n$, wherein $n = 1$ to 6.

35. The bioconjugate of claim 32, wherein *BAS* and *BAS** are different bioactive species.

36. A method of preparing the reagent of General Formula CIV comprising the step of condensing the reagent of General Formula CIII with a bioactive species to yield the reagent of General Formula CIV:



General Formula CIV

wherein group Z comprises a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with at least one of intermediate amide and disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length;

wherein group R_3 is selected from one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent; and


-131-

wherein *BAS* is a biologically active species.

37. The method of claim 36, wherein group R_3 is preferably selected from one of H, CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 , CH_2CH_2OH , and CH_2OCH_3 .

38. The method of claim 36, wherein group Z is preferably an unbranched alkyl chain of general formula $(CH_2)_n$, wherein $n = 1$ to 6.

39. A method of preparing the bioconjugate of General Formula CVI,



General Formula CVI

wherein group Z comprises a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with one of intermediate amide and disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length;

wherein group R_3 is selected is one of H, an alkyl, and a methylene or ethylene with an electronegative moiety;

wherein *BAS* is a biologically active species; and


wherein *BAS** is a biologically active species,

the method comprising the step of complexing the conjugate of claim 6 with a second conjugate comprising a second biologically active species and a phenylboronic acid.

40. The method of claim 39, wherein group R_3 is preferably selected from one of H, CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 , CH_2CH_2OH , and CH_2OCH_3 .

-132-

41. The method of claim 39, wherein group Z is preferably an unbranched alkyl chain of general formula $(CH_2)_n$, wherein $n = 1-6$.
42. The method of claim 39, wherein *BAS* and *BAS** are different bioactive species.
43. A method for preparing a conjugate of General Formula CIV:




General Formula CIV

wherein group Z comprises a spacer selected from a saturated or unsaturated chain of from about 0 to 6 carbon equivalents in length, an unbranched saturated or unsaturated chain of from about 6 to 18 carbon equivalents in length with one of intermediate amide and disulfide moieties, and a polyethylene glycol chain of from about 3 to 12 carbon equivalents in length;

wherein group R_3 is selected from one of H, an alkyl, and an alkyl or methylene with an electronegative moiety; and

wherein *BAS* is a biologically active species,
the method comprising:

condensing a reagent of general formula




with a hydroxylamine-derivative of the general formula NH_2OR_3 , wherein group R_2 is selected from one of an alkyl, and a methylene with an electronegative moiety.

-133-

44. The method of claim 43, wherein group R_3 is preferably selected from one of H, CH_3 , CH_2CH_3 , CH_2CN , CH_2COOH , CH_2CONH_2 , CH_2CH_2OH , and CH_2OCH_3 .


45. The method of claim 43, wherein group Z is preferably an unbranched alkyl chain of general formula $(CH_2)_n$, wherein $n = 1$ to 6.

46. The compound comprising:



wherein group R_3 is selected from one of H, an alkyl, and an alkyl or methylene with an electronegative moiety.


47. A method of making the compound of general formula:



wherein group R_3 is selected from one of H, an alkyl, and an alkyl or methylene with an electronegative moiety,


the method comprising:

condensing a methyl 2,6-dihydroxy-4-methylbenzoate of general formula:




-134-


wherein R_2 is an alkyl, with a suitable group Q to protect the amine moiety:




condensing the benzoate ester with suitable groups Q' to reduce the reactivity of the ring system:



hydrolyzing the benzoate ester to yield the salicylic acid:



condensing the acid with a hydroxylamine derivative of the general formula NH_2OR_3 , wherein R_3 is selected from one of an H, an alkyl, and a methylene or ethylene moiety with an electronegative substituent:



and removing the protecting groups Q and Q' .

1/14

Figure 1







Figure 2

Figure 3


4/14

Figure 4


Step 1




Step 2




Step 3



Step 4





Step 5




5/14


Figure 5



6/14




7/14








8/14

Figure 1'



9/14

*Figure 2'**Step 1**Step 2**Step 3**Step 4**Step 5*

10/14







*Figure 3'**Step 1**Step 2**Step 3**Step 4*

Figure 4'




12/14


Figure 5'



13/14



14/14



INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/US 97/13314

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07C233/49 G01N33/53 C07C259/10 C07F5/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 20591 A (PROLINX INC) 3 August 1995 see claims	1-47
A	US 3 293 252 A (J. FRIED EL AL) 20 December 1966 see the whole document	1-47

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"8" document member of the same patent family

Date of the actual completion of the international search

17 October 1997

Date of mailing of the international search report

30.10.97

Name and mailing address of the ISA

European Patent Office, P.B. 5616 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Sánchez García, J.M.

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/US 97/13314

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9520591 A	03-08-95	US 5594151 A	14-01-97
		US 5594111 A	14-01-97
		US 5623055 A	22-04-97
		AU 1732495 A	15-08-95
		CA 2181252 A	03-08-95
		CZ 9602201 A	11-12-96
		EP 0741734 A	13-11-96
		PL 315911 A	09-12-96
		SK 98996 A	05-02-97
		US 5648470 A	15-07-97
		US 5668257 A	16-09-97
		US 5668258 A	16-09-97
US 3293252 A	20-12-66	NONE	

Form PCT/ISA/210 (patent family annex) (July 1992)